

EE325 Introductory Econometrics (Section 1 semester 1/2020)

Assignment 4

Instruction: Write your answer in either paper or digital paper. However, if you write on paper, please scan it and save as a PDF file. Submission is via BE-Moodle as a PDF file for both cases. (Please keep the file below 10MB as that is the maximum per file capacity for student.)

Due date: Friday, November 6, 2020 (Before 10 P.M.)

1. From the data for 46 states in the United States for 1992, results of the regression are displayed as follows. $n = 46, k = 3$

$$\ln C_i = 4.30 - 1.34 \ln P_i + 0.17 \ln Y_i$$

$$se = (0.91) \quad (0.32) \quad (0.20)$$

$$\bar{R}^2 = 0.27$$

where C_i = cigarette consumption, packs per year

P_i = real price per pack, \$ per pack

Y_i = real disposable income per capita, \$ per week

1.1) Do the estimation results follow the law of demand?

1.2) What is the elasticity of demand for cigarettes with respect to price? Is it statistically significant? If so, is it statistically different from 1?

1.3) What is the income elasticity of demand for cigarettes? Is it statistically significant? If not, what might be the reasons for it?

2. From estimating the regression equation on net financial wealth (nettfa), age of the survey respondent (age), and annual family income (inc) for people in the United States. The wealth and income variables are both recorded in thousands of dollars. The OLS estimation results for the model are given by

$$nettfa_i = \beta_1 + \beta_2 inc_i + \beta_3 age_i + u_i$$

reg nettfa inc age

Sum square.

Source	SS	df	MS	Number of obs	=	9,275
Model	6414618.8	2	3207309.4	F(2, 9272)	=	943.21
Residual	31528770.7	9,272	3400.42825	Prob > F	=	0.0000
				R-squared	=	0.1691
				Adj R-squared	=	0.1689
Total	37943389.5	9,274	4091.3726	Root MSE	=	58.313

nettfa	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
inc	.9533566	.0252775	37.72	0.000	.9038072 1.002906
age	1.030777	.0591226	17.43	0.000	.9148838 1.14667
_cons	-60.69654	2.596333	-23.38	0.000	-65.78592 -55.60715

```
reg nettfafa inc age agesq
```

	Source	SS	df	MS	Number of obs	=	9,275
	Model	6567017.15	3	2189005.72	F(3, 9271)	=	646.80
	Residual	31376372.3	9,271	3384.35685	Prob > F	=	0.0000
	Total	37943389.5	9,274	4091.3726	R-squared	=	0.1731
					Adj R-squared	=	0.1728
					Root MSE	=	58.175

	nettfafa	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
	inc	.9782522	.0254891	38.38	0.000	.928288 1.028216
	age	-2.231489	.4897118	-4.56	0.000	-3.191432 -1.271547
	agesq	.0377221	.0056214	6.71	0.000	.026703 .0487413
	_cons	4.680388	10.08099	0.46	0.642	-15.08056 24.44134

2.1) Test the coefficient, in the first model, $\beta_3 < 1$ in the first model or not?

2.2) Due to estimation result by adding the age² variable or agesq. Perform the test whether we should include the quadratic term of the age variable or not? (Test for both t-test and F-test.) Also, interpret the meaning of this coefficient.

3. You are conducting an empirical investigation into the median prices of houses in 506 communities of a large metropolitan area. The sample data consist of 506 observations on the following observable variables:

P_i : the median house price in community i , in dollars;

NOX_i : the level of nitrous oxide in the air of community i , in parts per 100 million;

$DIST_i$: the weighted distance of community i from municipal area, in miles;

$ROOM_i$: the average number of rooms per house in community i ;

$STRAT_i$: the average student-teacher ratio of schools in community i .

Researcher estimates the following model of median house price. The OLS estimation results for the model are given by

$$\ln(P_i) = 11.08 - 0.9535 \ln(NOX_i) - 0.1343 \ln(DIST_i) + 0.2545 ROOM_i - 0.05245 STRAT_i$$

$$se = (0.3181) (0.1167) (0.04310) (0.01853) (0.005897)$$

$$RSS = 35.1835 \quad TSS = 84.5822$$

3.1) Interpret each of the coefficient estimates in regression equation.

3.2) Test the individual significance of each of the slope coefficient estimates for $\ln(NOX_i)$ and $ROOM_i$.

3.3) Find the R-squared, adjusted R-squared, and test the joint significance of all the slope coefficient estimates.

3.4) If researcher would like to test the proposition that the marginal effect of $\ln(NOX_i)$ on $\ln(P_i)$ equals the marginal effect of $\ln(DIST_i)$ on $\ln(P_i)$, write the restricted model and

perform the test comparing restricted and unrestricted model, given that OLS estimation of this restricted regression equation yields a Residual Sum of Squares value = 41.9532.

4. Production function (Y) of the industrial sector in Thailand. It depends on the capital factor (K) and labor factor (L) in the years 1980-2010 with the following estimation.

Model 1:

$$\ln Y_t = 18.27 + 0.536 \ln L_t + 0.024 \ln K_t$$

$$R^2 = 0.9389, RSS = 0.0124$$

Model 2:

$$\ln \left(\frac{Y}{L} \right)_t = 2.13 + 1.12 \ln \left(\frac{K}{L} \right)_t$$

$$R^2 = 0.8087, RSS = 0.0153$$

4.1) Interpret the coefficients of the independent variables in models 1 and 2.

4.2) Test the hypothesis. Is the industrial production function characterized by constant return to scale? (Hint: you can perform any type of test that you see fit.)

4.3) Can we compare the R^2 value between the two regression models? Why?

$$\therefore \beta_2 < 0$$

1

$$1.1) H_0: \beta_2 \geq 0$$


$$H_1: \beta_2 < 0$$

$$\alpha = 0.05$$

$$t_{cal} = \frac{\hat{\beta}_2 - \beta_2}{\text{se}(\hat{\beta}_2)} = \frac{-1.34 - 0}{0.32} = -4.1875$$

$$df = 46 - 3 = 43$$

critical value



Reject H_0 if $t_{cal} < -1.684$

And that $-4.1875 < -1.684$

\therefore Reject H_0

There is enough evidence to say that this result follows the law of demand at $\alpha = 0.05$

$$1.2) \epsilon_p = \frac{\% \Delta C}{\% \Delta P} = -1.34$$

$$1.3) \epsilon_y = \frac{\% \Delta C}{\% \Delta I} = 0.17$$

2.

$$2.1) \text{netff}_i = -0.696 + 0.9533 \text{inc}_i + 1.03077 \text{age}_i$$

$$\text{se} = (2.596) \quad (0.0252) \quad (0.591)$$

$$H_0: \beta_3 \geq 1$$

$$H_1: \beta_3 < 1$$

$$\alpha = 0.05$$

$$t_{cal} = \frac{\hat{\beta}_3 - \beta_3}{\text{se}(\hat{\beta}_3)} = \frac{1.03077 - 1}{0.591} = 0.052$$

$$df = n - k = 9275 - 3 = 9272$$

critical value



$$t_{0.05, 9272} = -1.645$$

β_3 is significant \therefore reject H_0

$\therefore \beta_3 \text{ is } \geq 1$

2.1) Test the coefficient, in the first model, $\rho_3 < 1$ in the first model or not.

2.2) Due to estimation result by adding the age² variable or agesq. Perform the test whether we should include the quadratic term of the age variable or not? (Test for both t-test and F-test.) Also, interpret the meaning of this coefficient.

F-test

$$neffq_i = \beta_1 + \beta_2 inc_i + \beta_3 age_i + \beta_4 age_i^2 + u_i \quad (\text{Unrestricted})$$

if $\beta_4 = 0$, we shouldn't include the term in

$\beta_4 \neq 0$, we should include in

$$H_0: \beta_4 = 0$$


$$H_1: \beta_4 \neq 0$$

$$\alpha = 0.05, N = n - k = 9 - 3 = 6$$

$$F_{ca} = \frac{RSS_R - RSS_{UR} / M}{RSS_{UR} / (n - k)} = \frac{R^2_{UR} - R^2_R / M}{(1 - R^2_{UR}) / (n - k)}$$
$$= \frac{(31528770.7 - 31376372.3) / 1}{31376372.3 / (9275 - 4)}$$

$$= 45.03024$$

Critical value



$$F_{0.05, (1, 9271)} = 3.89$$

reject H_0 , if $F_{ca} > 3.89$

Find that $45.03024 > 3.89$

\therefore reject H_0

There is enough evidence to say that the addition of age² to the model has significant, $\alpha = 5\%$.

3.1) Interpret each of the coefficient estimates in regression equation.

$B_1 = 11.08$; if $\ln(\text{NOX}_i)$, $\ln(\text{DIST}_i)$, ROOM_i , $\text{STRAT}_i = 0$, on average, the median house price will be equal to $e^{11.08} = 569840.88$

$B_2 = -0.9535$; if the level of nitrous oxide increase by 1%, on average, the median house price will decrease by 0.9535% holding other variable constant.

$B_3 = -0.1393$; if the weighted distance of community i increase by 1%, the median house price will decrease by 0.1393%.

$B_4 = 0.2545$; if the average number of rooms per house in community increase by 1%, the median house price will increase by 25.45%.

$B_5 = -0.05245$; if the average student-teacher ratio of school increase by 1%, on average, the median house price will decrease by 5.245% holding other variable constant.

3.2) Test the individual significance of each of the slope coefficient estimates for $\ln(\text{NOX}_i)$ and ROOM_i .

$\ln(\text{NOX}_i)$


$$H_0 : B_2 = 0$$

$$H_1 : B_2 \neq 0$$

$$t_{\text{cal}} = \frac{\hat{B}_2 - B_2}{\text{se}(\hat{B}_2)} = \frac{-0.9535 - 0}{0.1767} = -5.4012$$

$$d.f = n - k = 506 - 5 = 501 = \infty$$

Critical value



reject H_0
if $t_{\text{cal}} < -1.96$ or $t_{\text{cal}} > 1.96$

And that $-5.4012 < -1.96$

\therefore Reject H_0

There is enough evidence to say that $\ln(\text{NOX})$ has significant at $\alpha = 0.05$

$$H_0 : B_4 = 0$$

$$H_1 : B_4 \neq 0$$


$$\alpha = 0.05$$

$$t_{\text{cal}} = \frac{\hat{B}_4 - B_4}{\text{se}(\hat{B}_4)} = \frac{0.2545 - 0}{0.01853}$$

$$= 13.73948$$

$$d.f = n - k = 506 - 5 = 501 = \infty$$

critical value



\therefore reject H_0

$\therefore B_4$ is significant

$\ln(NOX_i)$ and $ROOM_i$.

overall

3.3) Find the R-squared, adjusted R-squared, and test the joint significance of all the slope coefficient estimates.

$$H_0: \beta_2 = \beta_3 = \beta_4 = \beta_5 = 0$$

H_1 : Not all slope coefficient are simultaneously zero

$$\alpha = 0.05$$

$$ESS = TSS - RSS = 89.5822 - 35.1835 = 49.3987$$

$$F_{ca} = \frac{ESS/k-1}{RSS/n-k} = \frac{R^2/k-1}{(1-R^2)/n-k}$$

$$= \frac{49.3987/4}{35.1835/501} = 175.8991$$

reject $H_0 \rightarrow$ This is enough evidence to say that at least one parameter is not equal to zero at $\alpha = 5\%$.



$$F_{0.05}(4, 501) = 2.37$$

coefficient estimates.

3.4) If researcher would like to test the proposition that the marginal effect of $\ln(NOX_i)$ on $\ln(P_i)$ equals the marginal effect of $\ln(DIST_i)$ on $\ln(P_i)$, write the restricted model and

$$H_0: \beta_2 = \beta_3 \text{ or } \beta_2 - \beta_3 = 0$$

$$H_1: \beta_2 \neq \beta_3 \text{ or } \beta_2 - \beta_3 \neq 0$$

$$\alpha = 0.05$$


$$\ln(P_i) = \beta_1 + \beta_2 \ln(NOX_i) + \beta_3 \ln(DIST_i) + \beta_4 ROOM_i + \beta_5 STRAT_i + u_i$$

$$= \beta_1 + \beta_2 \ln(NOX_i) + \beta_2 \ln(DIST_i) + \beta_4 ROOM_i + \beta_5 STRAT_i + u_i$$

$$= \beta_1 + \beta_2 \ln(NOX_i) + \ln(DIST_i) + \beta_4 ROOM_i + \beta_5 STRAT_i + u_i \text{ (restricted)}$$

$$F_{ca} = \frac{RSS_R - RSS_{UR}}{m} \cdot \frac{n-k}{RSS_{UR}}$$

$$= \frac{41.9532 - 35.1835/1}{35.1835/504-5} = 96.39801$$



$$F_{0.05}(1, 501) = 3.84$$

\therefore reject $H_0 \therefore \beta_2 \neq \beta_3$

4.1) Interpret the coefficients of the independent variables in models 1 and 2.

$$\text{model 1: } \ln Y_t = \beta_1 + \beta_2 \ln l_t + \beta_3 \ln k_t + u_t = \text{unrestricted}$$

$$\text{model 2: } \ln Y_t = \beta_1 + \beta_3 \ln \left(\frac{k}{l} \right) + u_t \quad (\text{restricted})$$

$$\text{condition is } \beta_2 + \beta_3 = 1$$

$$H_0: \beta_2 + \beta_3 = 1$$

$$H_1: \beta_2 + \beta_3 \neq 1$$

$$\alpha = 0.05$$

$$F = \frac{(R_{UR}^2 - R_R^2) / m}{(1 - R_{UR}^2) / (n - k_{UR})} = \frac{(0.9389 - 0.8087) / 1}{(1 - 0.9389) / 27} = 57.535$$

find critical value $F_{0.05, 1, 27} \approx 4.23$

reject H_0 when $F > F_{0.05, 1, 27}$ because $F = 57.535 > F_{0.05, 1, 27} = 4.23$

\therefore Reject H_0

4.3) Can we compare the R^2 value between the two regression models? Why?

we cannot compare R^2 value between two regression models because it has different variables. since in model 1 has $\ln Y_t$ while model 2 has $\left(\frac{Y}{L} \right)_t$.