

AUTOCORRELATION

EE 325 (Ajarn Kaewkwan Tangtipongkul)

THE NATURE OF THE PROBLEM

EE 325 (Ajarn Kaewkwan Tangtipongkul)

THE NATURE OF THE PROBLEM

Autocorrelation may be defined as “correlation between members of series of observations ordered in time (as in time series data) or space (as in cross-sectional data)”

$$\text{cov}(u_i, u_j | x_i, x_j) = E(u_i u_j) = 0 \quad i \neq j$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

Autocorrelation defines as Lag correlation of a given series with itself, lagged by a number of time units

The correlation between two time series such as $u_1, u_2, \dots, u_{10}$ and $u_2, u_3, \dots, u_{11}$, where the former is the latter series lagged by one time period

EE 325 (Ajarn Kaewkwan Tangtipongkul)

Serial Correlation is the correlation between time series such as $u_1, u_2, \dots, u_{10}$ and $v_2, v_3, \dots, v_{11}$, where u and v are two different time series

EE 325 (Ajarn Kaewkwan Tangtipongkul)

NO AUTOCORRELATION

$$E(u_i u_j) = 0, \quad i \neq j$$

$$E(uu') = \begin{bmatrix} E(u_1^2) & E(u_1 u_2) & \dots & E(u_1 u_n) \\ E(u_2 u_1) & E(u_2^2) & \dots & E(u_2 u_n) \\ \dots & \dots & \dots & \dots \\ E(u_n u_1) & E(u_n u_2) & \dots & E(u_n^2) \end{bmatrix}$$

$$E(uu') = \begin{bmatrix} \sigma^2 & 0 & 0 & 0 \\ 0 & \sigma^2 & 0 & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \sigma^2 \end{bmatrix} = \sigma^2 \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 1 \end{bmatrix} = \sigma^2 I$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

WHY DOES SERIAL CORRELATION OCCUR?

EE 325 (Ajarn Kaewkwan Tangtipongkul)

INERTIA

A salient feature of most economic time series is sluggishness


Example

GNP, price indexes, production, employment, and unemployment exhibit business cycles

EE 325 (Ajarn Kaewkwan Tangtipongkul)

- Specification Bias: excluded variables case

EE 325 (Ajarn Kaewkwan Tangtipongkul)



$$Y_t = \beta_1 + \beta_2 X_{2t} + \beta_3 X_{3t} + \beta_4 X_{4t} + u_t$$

Y_t = quantity of beef demand

X_2 = price of beef

X_3 = consumer income

X_4 = price of pork

t = time

$$Y_t = \beta_1 + \beta_2 X_{2t} + \beta_3 X_{3t} + v_t$$

$$v_t = \beta_4 X_{4t} + u_t$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

The price of pork affects the consumption of beef, the error or disturbance term V will reflect a systematic pattern, thus creating (false) autocorrelation

EE 325 (Ajarn Kaewkwan Tangtipongkul)

SPECIFICATION BIAS: INCORRECT FUNCTIONAL FORM

Suppose the true or correct model in a cost-output study is as follows:

$$MARGINAL\ COST_i = \beta_1 + \beta_2 output_i + \beta_3 output_i^2 + u_i$$


But we fit the following model:

$$MARGINAL\ COST_i = \alpha_1 + \alpha_2 output_i + v_i$$

$$v_i = \beta_3 output_i^2 + u_i$$

The marginal cost curve corresponding to the true model is shown in the next figure along with the incorrect linear cost curve

EE 325 (Ajarn Kaewkwan Tangtipongkul)



EE 325 (Ajarn Kaewkwan Tangtipongkul)

COBWEB PHENOMENON

The supply of many agricultural commodities reflects the so-called cobweb phenomenon, where supply reacts to price with a lag of one time period because supply decisions take time to implement.

$$Supply_t = \beta_1 + \beta_2 P_{t-1} + u_t$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

LAGS

$$Consumption_t = \beta_1 + \beta_2 income + \beta_3 consumption_{t-1} + u_t$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

Data Transformation

$$Y_t = \beta_1 + \beta_2 X_t + u_t$$

$$Y = Consumption \quad Expenditure$$

$$X = income$$

$$Y_{t-1} = \beta_1 + \beta_2 X_{t-1} + u_{t-1} \quad \leftarrow \text{Level Form}$$

$$\Delta Y_t = \beta_2 \Delta X_t + \Delta u_t$$

$$\Delta Y_t = \beta_2 \Delta X_t + v_t$$

$$v_t = \Delta u_t = (u_t - u_{t-1}) \quad \leftarrow \text{Difference Form}$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

$$v_t = u_t - u_{t-1}$$


$$E(u) = 0$$

$$E(v_t) = E(u_t - u_{t-1}) = E(u_t) - E(u_{t-1}) = 0$$

$$\text{var}(v_t) = \text{var}(u_t - u_{t-1}) = \text{var}(u_t) - \text{var}(u_{t-1}) = 2\sigma^2$$

$$\text{cov}(v_t, v_{t-1}) = E(v_t v_{t-1}) = E[(u_t - u_{t-1})(u_{t-1} - u_{t-2})] = -\sigma^2$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)



OLS ESTIMATION IN THE PRESENCE OF AUTOCORRELATION

EE 325 (Ajarn Kaewkwan Tangtipongkul)

$$Y_t = \beta_1 + \beta_2 X_t + u_t$$

$$u_t = \rho u_{t-1} + \varepsilon_t \quad -1 < \rho < 1$$

ρ (coefficient of autocovariance)

$$E(\varepsilon_t) = 0$$

$$\text{var}(\varepsilon_t) = \sigma_\varepsilon^2$$

$$\text{cov}(\varepsilon_t, \varepsilon_{t+s}) = 0 \quad s \neq 0$$

ε (white noise error term)

EE 325 (Ajarn Kaewkwan Tangtipongkul)

$$u_t = \rho u_{t-1} + \varepsilon_t \quad -1 < \rho < 1$$

This equation is known as a Markov first order autoregressive scheme or first-order autoregressive scheme AR(1)

EE 325 (Ajarn Kaewkwan Tangtipongkul)

AR (1)

$$E(u_t) = \rho E(u_{t-1}) + E(\varepsilon) = 0$$

$$\text{var}(u_t) = E(u_t^2)$$

$$= \rho^2 \text{var}(u_{t-1}) + \text{var}(\varepsilon) = \frac{\sigma_\varepsilon^2}{1 - \rho^2}$$

$$\text{var}(u_t) = \text{var}(u_{t-1}) = \sigma^2$$

$$\text{var}(\varepsilon_t) = \sigma_\varepsilon^2$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

$$\text{cov}(u_t, u_{t+s}) = E(u_t, u_{t+s}) = \rho^s \frac{\sigma_\varepsilon^2}{1 - \rho^2}$$

Covariance between error terms s periods apart

The symmetry property of covariances

$$\text{cov}(u_t, u_{t+s}) = \text{cov}(u_t, u_{t-s})$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

- By definition, the (population) coefficient of correlation between u_t and u_{t-1} is

$$\rho = \frac{E\{[u_t - E(u_t)][u_{t-1} - E(u_{t-1})]\}}{\sqrt{\text{var}(u_t)}\sqrt{\text{var}(u_{t-1})}} = \frac{E(u_t u_{t-1})}{\text{var}(u_{t-1})}$$

Since $E(u_t) = 0$ for each t and $\text{var}(u_t) = \text{var}(u_{t-1})$ because we are retaining the assumption of homoscedasticity.

EE 325 (Ajarn Kaewkwan Tangtipongkul)

$$\text{Cor}(u_t, u_{t+s}) = \rho^s$$

Correlation between error terms s period apart

The symmetry property of correlations

$$\text{Cor}(u_t, u_{t+s}) = \text{Cor}(u_t, u_{t-s})$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

Since ρ is a constant between -1 and +1, , variance equation shows that under the AR(1) scheme, the variance of u_t is still homoscedastic, but u_t is correlated not only with its immediate past value but its values several periods in the past.

EE 325 (Ajarn Kaewkwan Tangtipongkul)

Two variable regression model

$$Y_t = \beta_1 + \beta_2 X_t + u_t$$

OLS estimator of the slope coefficient is

$$\hat{\beta}_2 = \frac{\sum (X_t - \bar{X})(Y_t - \bar{Y})}{\sum (X_t - \bar{X})^2} = \frac{\sum x_t y_t}{\sum x_t^2}$$

and its variance is given by

$$\text{var}(\hat{\beta}_2) = \frac{\sigma^2}{\sum (X_t - \bar{X})^2} = \frac{\sigma^2}{\sum x_t^2}$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

Now under the AR(1) scheme, it can be shown that the variance of this estimator is

$$\text{var}(\hat{\beta}_2)_{AR1} = \frac{\sigma^2}{\sum x_t^2} \left[1 + 2\rho \frac{\sum x_t x_{t-1}}{\sum x_t^2} + 2\rho^2 \frac{\sum x_t x_{t-2}}{\sum x_t^2} + \dots + 2\rho^{n-1} \frac{\sum x_t x_n}{\sum x_t^2} \right]$$

where

$$x_t = (X_t - \bar{X})$$

$$x_{t-1} = (X_{t-1} - \bar{X})$$

$\vdots$

$$x_n = (X_n - \bar{X})$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

The difference between the $\text{var}(\hat{\beta}_2)$ and $\text{var}(\hat{\beta}_2)_{AR(1)}$, assume that the regressor X also follows the first order autoregressive scheme with a coefficient of autocorrelation of r

$$\text{var}(\hat{\beta}_2)_{AR(1)} = \frac{\sigma^2}{\sum x_t^2} \left(\frac{1+r\rho}{1-r\rho} \right) = \text{var}(\hat{\beta}_2)_{OLS} \left(\frac{1+r\rho}{1-r\rho} \right)$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

As in the case of **heteroscedasticity** in the presence of **autocorrelation** the OLS estimators are still linear unbiased as well as consistent and asymptotically normally distributed, but they are **no longer efficient** (i.e., **minimum variance**)


EE 325 (Ajarn Kaewkwan Tangtipongkul)

CONSEQUENCES OF USING OLS IN THE PRESENCE OF AUTOCORRELATION

EE 325 (Ajarn Kaewkwan Tangtipongkul)

OLS ESTIMATION ALLOWING FOR AUTOCORRELATION

$\hat{\beta}_2$ is not BLUE and even if we use $\text{var}(\hat{\beta}_2)_{AR1}$, the confidence intervals derived from there are likely to be wider than those based on the GLS procedure



OLS ESTIMATION DISREGARDING AUTOCORRELATION

The situation is potentially very serious if we not only use $\hat{\beta}_2$ but also continue to use $\text{var}(\hat{\beta}_2) = \sigma^2 / \sum (X_i - \bar{X})^2$, which completely disregards the problem of autocorrelation, that is, we mistakenly believe that the usual assumptions of the classical model hold true. Errors will arise for the following reasons

EE 325 (Ajarn Kaewkwan Tangtipongkul)

1. The residual variance $\hat{\sigma}^2 = \frac{\sum \hat{u}_i^2}{(n-2)}$ is likely to underestimate the true σ^2
2. As a result, we are likely to overestimate R^2
3. Even if σ^2 is not underestimated, $\text{var}(\hat{\beta}_2)$ may underestimated $\text{var}(\hat{\beta}_2)_{AR1}$ its variance under AR(1), even though the latter is inefficient compared to $\text{var}(\hat{\beta}_2)^{GLS}$
4. Therefore, the usual t and F tests of significance are no longer valid

EE 325 (Ajarn Kaewkwan Tangtipongkul)

EXAMPLE

To see how OLS is likely to underestimate σ^2 and the variance of $\hat{\beta}_2$, let us conduct the following **Monte Carlo experiment**

Suppose in the two-variable model

$$Y_t = 1.0 + 0.8X_t + u_t$$

$$E(Y_t | X_t) = 1.0 + 0.8X_t$$

$$u_t = 0.7u_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim N(0,1)$$


EE 325 (Ajarn Kaewkwan Tangtipongkul)

TABLE 12.1
A Hypothetical
Example of Positively
Autocorrelated Error
Terms

	e_t	$u_t = 0.7u_{t-1} + e_t$
0	0	$u_0 = 5$ (assumed)
1	0.464	$u_1 = 0.7(5) + 0.464 = 3.964$
2	2.026	$u_2 = 0.7(3.964) + 2.0262 = 4.8008$
3	2.455	$u_3 = 0.7(4.8010) + 2.455 = 5.8157$
4	-0.323	$u_4 = 0.7(5.8157) - 0.323 = 3.7480$
5	-0.068	$u_5 = 0.7(3.7480) - 0.068 = 2.5556$
6	0.296	$u_6 = 0.7(2.5556) + 0.296 = 2.0849$
7	-0.288	$u_7 = 0.7(2.0849) - 0.288 = 1.1714$
8	1.298	$u_8 = 0.7(1.1714) + 1.298 = 2.1180$
9	0.241	$u_9 = 0.7(2.1180) + 0.241 = 1.7236$
10	-0.957	$u_{10} = 0.7(1.7236) - 0.957 = 0.2495$

Note: e_t data obtained from *A Million Random Digits and One Hundred Thousand Deviates*, Rand Corporation, Santa Monica, Calif., 1950.

EE 325 (Ajarn Kaewkwan
Tangtipongkul)



EE 325 (Ajarn Kaewkwan
Tangtipongkul)

Now suppose the values of X are fixed at
1,2,3,...,10.

$$\hat{Y}_t = 6.5452 + 0.3051X_t$$


(0.6153) (0.0992)

$$t = (10.6366) (3.0763)$$

$$r^2 = 0.5419$$

$$\hat{\sigma}^2 = 0.8114$$

EE 325 (Ajarn Kaewkwan
Tangtipongkul)



EE 325 (Ajarn Kaewkwan
Tangtipongkul)

TABLE 12.2
Generation of Y
Sample Values

X_t	u_t	$Y_t = 1.0 + 0.8X_t + u_t$
1	3.9640	$Y_1 = 1.0 + 0.8(1) + 3.9640 = 5.7640$
2	4.8010	$Y_2 = 1.0 + 0.8(2) + 4.8008 = 7.4008$
3	5.8157	$Y_3 = 1.0 + 0.8(3) + 5.8157 = 9.2157$
4	3.7480	$Y_4 = 1.0 + 0.8(4) + 3.7480 = 7.9480$
5	2.5556	$Y_5 = 1.0 + 0.8(5) + 2.5556 = 7.5556$
6	2.0849	$Y_6 = 1.0 + 0.8(6) + 2.0849 = 7.8849$
7	1.1714	$Y_7 = 1.0 + 0.8(7) + 1.1714 = 7.7714$
8	2.1180	$Y_8 = 1.0 + 0.8(8) + 2.1180 = 9.5180$
9	1.7236	$Y_9 = 1.0 + 0.8(9) + 1.7236 = 9.9236$
10	0.2495	$Y_{10} = 1.0 + 0.8(10) + 0.2495 = 9.2495$

Note: u_t data obtained from Table 12.1.

EE 325 (Ajarn Kaewkwan
Tangtipongkul)

Keeping the X_t and ε_t given in Table 2.1
and Table 2.2, let us assume $\rho = 0$, that
is, no autocorrelation

TABLE 12.3
Sample of Y Values
with Zero Serial
Correlation

X_t	$e_t = u_t$	$Y_t = 1.0 + 0.8X_t + e_t$
1	0.464	2.264
2	2.026	4.626
3	2.455	5.855
4	-0.323	3.877
5	-0.068	4.932
6	0.296	6.096
7	-0.288	6.312
8	1.298	8.698
9	0.241	8.441
10	-0.957	8.043

Note: Since there is no autocorrelation, the u_t and e_t are identical. The e_t are from Table 12.1.

EE 325 (Ajarn Kaewkwan
Tangtipongkul)

THE REGRESSION BASED ON TABLE 12.3 IS AS FOLLOWS:

$$\hat{Y}_t = 2.5345 + 0.6145X_t$$

(0.6796) (0.1087)

$$t = (3.7910) (5.6541)$$

$$r^2 = 0.7997$$

$$\hat{\sigma}^2 = 0.9752$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

DETECTING AUTOCORRELATION

- Durbin-Watson d Test

EE 325 (Ajarn Kaewkwan Tangtipongkul)

DURBIN-WATSON D TEST

$$d = \frac{\sum_{t=2}^{t=n} (\hat{u}_t - \hat{u}_{t-1})^2}{\sum_{t=1}^{t=n} \hat{u}_t^2}$$

The ratio of the sum of squared differences in successive residuals to the RSS. Note that in the numerator of the d statistic the number of observations is n-1 because one observation is lost in taking successive differences

EE 325 (Ajarn Kaewkwan Tangtipongkul)

ASSUMPTIONS UNDERLYING THE D STATISTIC

- The regression model includes the intercept term
- The explanatory variables, the X's are nonstochastic, or fixed in repeated sampling
- The disturbances are generated by the AR(1)

$$u_t = \rho u_{t-1} + \varepsilon_t$$

Therefore, it cannot be used to detect higher-order AR schemes

- The error term u_t is assumed to be normally distributed

EE 325 (Ajarn Kaewkwan Tangtipongkul)

- The regression model does not include the lagged value(s) of the dependent variable as one of the explanatory variables
- There are no missing observations in the data

EE 325 (Ajarn Kaewkwan Tangtipongkul)

$$d = \frac{\sum_{t=2}^{t=n} (\hat{u}_t - \hat{u}_{t-1})^2}{\sum_{t=1}^{t=n} \hat{u}_t^2}$$

$$d = \frac{\sum \hat{u}_t^2 + \sum \hat{u}_{t-1}^2 - 2 \sum \hat{u}_t \hat{u}_{t-1}}{\sum \hat{u}_t^2}$$

$$d \approx 2 \left(1 - \frac{\sum \hat{u}_t \hat{u}_{t-1}}{\sum \hat{u}_t^2} \right)$$

Since $\sum \hat{u}_t^2$ and $\sum \hat{u}_{t-1}^2$ differ in only one observation, they are approximately equal. Therefore, setting

$$\sum \hat{u}_{t-1}^2 \approx \sum \hat{u}_t^2$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

$$\hat{\rho} = \frac{\sum \hat{u}_t \hat{u}_{t-1}}{\sum \hat{u}_t^2}$$

$$d \approx 2(1 - \hat{\rho})$$

But since $-1 \leq \rho \leq 1$, $0 \leq d \leq 4$


EE 325 (Ajarn Kaewkwan Tangtipongkul)

The mechanics of the Durbin-Watson test are

1. Run the OLS regression and obtain the residuals d_i
2. Compute d
3. For the given sample size and given number of explanatory variables, find out the critical and d_U values
4. Now follow the decision rules given in Table 12.6

EE 325 (Ajarn Kaewkwan Tangtipongkul)

DURBIN-WATSON D STATISTIC



EE 325 (Ajarn Kaewkwan Tangtipongkul)

TABLE 12.6
Durbin-Watson d
Test: Decision Rules

Null Hypothesis	Decision	If
No positive autocorrelation	Reject	$0 < d < d_L$
No positive autocorrelation	No decision	$d_L \leq d \leq d_U$
No negative correlation	Reject	$4 - d_L < d < 4$
No negative correlation	No decision	$4 - d_U \leq d \leq 4 - d_L$
No autocorrelation, positive or negative	Do not reject	$d_U < d < 4 - d_U$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

EXAMPLE

- U.S. Consumption Expenditure for the period 1947-2000

$$\ln Consumption = \beta_1 + \beta_2 \ln income + \beta_3 \ln wealth + \beta_4 Interest$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

$$\ln Consumption = -0.4677 + 0.8049 \ln income + 0.2013 \ln wealth + 0.0027 Interest$$

Source	SS	df	MS	Number of obs =	54
Model	16.1637474	3	5.3879158	F(3, 50) =	37832.86
Residual	.007120721	50	.000142414	Prob > F =	0.0000
Total	16.1708681	53	.305110719	R-squared =	0.9996
				Adj R-squared =	0.9995
				Root MSE =	.01193

	Inc	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
lnincome		.8048728	.0174978	46.00	0.000	.7697273 .8400182
lnwealth		.2012702	.0175926	11.44	0.000	.1659345 .236606
i		-.0026891	.0007619	-3.53	0.001	-.0042194 -.0011587
_cons		-.467712	.042778	-10.93	0.000	-.5536342 -.3817899


EE 325 (Ajarn Kaewkwan Tangtipongkul)

HYPOTHESIS TESTING

H_0 : No positive autocorrelation
 H_0' : No negative autocorrelation
 H_1 : otherwise
 H_1' : otherwise

EE 325 (Ajarn Kaewkwan Tangtipongkul)

DURBIN-WATSON D STATISTIC



EE 325 (Ajarn Kaewkwan Tangtipongkul)

DURBIN-WATSON TEST

```

. tsset year
  time variable: year, 1947 to 2000
    delta: 1 unit

. estat dwatson
Durbin-Watson d-statistic( 4, 54) = 1.289232

(n = 54, k = 4) At 5% Significance level
d_L = 1.414
d_U = 1.724
Reject null hypothesis
Evidence of positive autocorrelation
    
```

EE 325 (Ajarn Kaewkwan Tangtipongkul)

CORRECTING FOR (PURE) AUTOCORRELATION

EE 325 (Ajarn Kaewkwan Tangtipongkul)

CORRECTING FOR (PURE) AUTOCORRELATION

1. The method of Generalized Least Squares (GLS) method
2. The Newey West method – to obtain standard errors of OLS estimators that are corrected for autocorrelation

EE 325 (Ajarn Kaewkwan Tangtipongkul)

THE METHOD OF GENERALIZED LEAST SQUARES (GLS)

The two-variable regression model:

$$Y_t = \beta_1 + \beta_2 X_t + u_t$$

and assume that the error term follows the AR(1) scheme, namely,

$$u_t = \rho u_{t-1} + \varepsilon_t, \quad -1 < \rho < 1$$

Now we consider two cases:

- (1) ρ is known
- (2) ρ is not known but has to be estimated

EE 325 (Ajarn Kaewkwan Tangtipongkul)

When ρ is known

$$Y_{t-1} = \beta_1 + \beta_2 X_{t-1} + u_{t-1}$$

$$\rho Y_{t-1} = \rho \beta_1 + \rho \beta_2 X_{t-1} + \rho u_{t-1}$$

$$(Y_t - \rho Y_{t-1}) = \beta_1(1 - \rho) + \beta_2(X_t - \rho X_{t-1}) + (u_t - \rho u_{t-1})$$

$$(Y_t - \rho Y_{t-1}) = \beta_1(1 - \rho) + \beta_2(X_t - \rho X_{t-1}) + \varepsilon_t$$

$$\text{where } \varepsilon_t = (u_t - \rho u_{t-1})$$

$$Y_t^* = \beta_1^* + \beta_2^* X_t^* + \varepsilon_t$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

Since the error term in

$$Y_t^* = \beta_1^* + \beta_2^* X_t^* + \varepsilon_t$$

satisfies the usual OLS assumptions, we can apply OLS to the transformed variables X^* and Y^* and obtain estimators with all the optimum properties, namely, BLUE.

EE 325 (Ajarn Kaewkwan Tangtipongkul)

When ρ is not known

- based on Durbin-Watson d Statistic

EE 325 (Ajarn Kaewkwan Tangtipongkul)

ρ BASED ON DURBIN-WATSON D STATISTIC

$$\hat{\rho} \approx 1 - \frac{d}{2}$$

In reasonably large sample one can obtain ρ and use it to transform the data as shown in the generalized difference equation

$$(Y_t - \rho Y_{t-1}) = \beta_1(1 - \rho) + \beta_2(X_t - \rho X_{t-1}) + \varepsilon_t$$

$$\text{where } \varepsilon_t = (u_t - \rho u_{t-1})$$

EE 325 (Ajarn Kaewkwan Tangtipongkul)

THE NEWEY WEST METHOD

EE 325 (Ajarn Kaewkwan Tangtipongkul)

THE NEWEY WEST METHOD OF CORRECTING THE OLS STANDARD ERRORS

The corrected standard errors are known as **HAC (Heteroscedasticity and autocorrelation-consistent) standard errors** or simply **Newey West standard errors**

EE 325 (Ajarn Kaewkwan Tangtipongkul)

THE NEWEY WEST METHOD

```
. newey lnc lnincome lnwealth i, lag(3)
```

Regression with Newey-West standard errors
maximum lag: **3**

Number of obs = **54**
F(3, 50) = **22341.29**
Prob > F = **0.0000**

lnc	Coef.	Newey-West Std. Err.	t	P> t	[95% Conf. Interval]	
lnincome	.8048728	.0171172	47.02	0.000	.7704919	.8392536
lnwealth	.2012702	.0154469	13.03	0.000	.1702441	.2322963
i	-.0020891	.0008798	-3.06	0.004	-.0044563	-.0009219
_cons	-.467712	.0439367	-10.65	0.000	-.5559616	-.3794625

EE 325 (Ajarn Kaewkwan
Tangtipongkul)

SOURCE

Gujarati, D.N. (2009) Basic Econometrics. 5th ed.
Singapore, McGraw-Hill.

EE 325 (Ajarn Kaewkwan
Tangtipongkul)