

5 Exponential and Logarithmic Functions

5.1 Exponential Functions (Revision)



Exponential functions are used extensively in modelling and solving a while variety of real-world problems, including growth of money in compound interest; growth of populations of people, animals, and bacteria; radioactive decay; and learning associated with new technology e.g. computing and manufacturing technology.

5.1.1 General form:

$$y = f(x) = b^x \quad b > 0, b \neq 1$$

Domain is $\mathbb{R}$ and range is $\mathbb{R}^+$.

b is positive to avoid complex (imaginary) number.




important features of exponential function

- All graphs will pass through the point $(0, 1)$.
- All graphs are continuous.
- The x axis is a horizontal asymptote.
- If $b > 1$, then b^x increases as x increases.
- If $0 < b < 1$, then b^x decreases as x decreases.

typical graph shape
(The shape depends on values of b)

5.1.2 Base e Exponential Function

$$e = 2.718\ 281\ 828\ 459\ \dots$$



→ similar to graphs of $y = b^x$

again the shape depends on value of b , in this case for $y = e^{-x}$

$$\Rightarrow y = \left(\frac{1}{e}\right)^x$$

$\therefore b = \frac{1}{e}$ which is < 1

5.1.3 Exponential Properties

If n and m are positive integers,

$$b^{\frac{n}{m}} = (\sqrt[m]{b})^n = \sqrt[m]{b^n}$$

where $\sqrt[m]{b}$ denotes the positive m th root

$$b^{-n} = \frac{1}{b^n}$$

$$b^0 = 1$$

$$b^x = b^y \text{ if and only if } x = y$$

$$b^m b^n = b^{m+n}$$

$$(b^n)^m = b^{mn}$$

$$(ab)^n = a^n b^n$$

$$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}$$

$$\frac{a^m}{a^n} = a^{m-n} = \frac{1}{a^{n-m}}$$

On the other hand, $y = e^x$
 $b = e > 1$

→ please learn this!!



5.2 Logarithmic Functions (Revision)

inverse $\Rightarrow$ exponential
in function e.g.

Logarithmic function is inversed of exponential function.

5.2.1 General form:

$$y = \log_b x \quad \text{where} \quad x = b^y, b > 0, b \neq 1$$

Domain is $\mathbb{R}^+$ and range is $\mathbb{R}$.

* Note that Domain of log = Range of exponential
Range Domain

Note that $\log_b 1 = 0$. If the base b is 10, it is normally omitted. The natural logarithmic of

x , $\ln(x)$ is $\log_e(x)$. (natural log $\therefore$ find in real life examples often)

Note that y of
log functions = x
of exponential

and x of log =
 y of exponential




FIGURE 2

Exponential Function		Logarithmic Function	
x	$y = 2^x$	$x = 2^y$	y
-3	$\frac{1}{8}$	$\frac{1}{8}$	-3
-2	$\frac{1}{4}$	$\frac{1}{4}$	-2
-1	$\frac{1}{2}$	$\frac{1}{2}$	-1
0	1	1	0
1	2	2	1
2	4	4	2
3	8	8	3

Ordered
pairs
reversed

$\hookrightarrow$ note the shape $y = \log_2 x$ is symmetrical with $y = 2^x$

5.2.2 Logarithmic Properties ($b > 0$ and $b \neq 1$, u and v are also positive numbers)

$$\log_b 1 = 0 \quad \text{and} \quad \log_b b = 1$$

$$\log_b u = \log_b v \quad \text{if and only if} \quad u = v$$

$$\log_b (uv) = \log_b u + \log_b v$$

$$\log_b \left(\frac{u}{v} \right) = \log_b u - \log_b v$$

$$* \log_b u^r = r \log_b u$$

$$* \log_b b^u = u$$

$$* \log_b x = \frac{\ln x}{\ln b}$$

learn this !!

the line $y = x$
as axis of
symmetrical

* are often in differentiation & integration of log functions

5.3 Differentiation of Logarithmic and Exponential Functions

① $\frac{d}{dx}(e^x) = e^x$	② $\frac{d}{dx}(e^u) = e^u \frac{du}{dx}$
③ $\frac{d}{dx}(\ln x) = \frac{1}{x}$	④ $\frac{d}{dx}(\ln u) = \frac{1}{u} \frac{du}{dx}$

I'll give these in exam but you have to learn how to use these

Ex. 1: Find the derivative of the following functions.

(solution in separate files)

(a) $y = 2e^x$

Ans: $y' = 2e^x$

(b) $y = 2e^{3x}$

Ans: $y' = 6e^{3x}$

(c) $y = ae^{-bt}$

Ans: $y' = -abe^{-bt}$

(d) $y = 5e^{x^4+2x}$

Ans: $y' = 5(4x^3 + 2)e^{x^4+2x}$

(e) $y = \frac{e^{-3x}}{x^2 + 1}$

Ans: $y' = e^{-3x} \left[\frac{-3x^2 - 2x - 3}{(x^2 + 1)^2} \right]$

Ex. 2: Show that the derivative of $f(x) = xe^{2x}$ is $f'(x) = (2x+1)e^{2x}$. In addition, find the largest and the smallest values of the function $f(x) = xe^{2x}$ on the interval $-1 \leq x \leq 1$.

Ans: Minimum $f\left(-\frac{1}{2}\right) \approx -0.184$ and maximum $f(1) \approx 7.389$ (solution in separate file)

Ex. 3: Find the derivative of the following functions.

(solution in separate file)

(a) $y = 5 \ln x$

Ans: $y' = \frac{5}{x}$

(b) $y = \ln(5x)$

Ans: $y' = \frac{1}{x}$

(c) $y = x \ln x$

Ans: $y' = 1 + \ln x$

$$(d) y = \frac{\ln \sqrt[3]{x^2}}{x^4}$$

$$\text{Ans: } y' = \frac{2}{3} \left[\frac{1 - 4 \ln x}{x^5} \right]$$

$$(e) y = 3 \ln(x^2 - 5x)$$

$$\text{Ans: } y' = \frac{3(2x - 5)}{x^2 - 5x}$$

$$(f) y = (x + \ln x)^{\frac{3}{2}}$$

$$\text{Ans: } y' = \frac{3}{2} (x + \ln x)^{\frac{1}{2}} \left(1 + \frac{1}{x} \right)$$

$$(g) y = \log_b x$$

$$\text{Ans: } y' = \frac{1}{(\ln b)x}$$

$$(h) y = \log_b u \text{ (} u \text{ is a function of } x \text{)}$$

$$\text{Ans: } y' = \frac{1}{(\ln b)u} \cdot \frac{du}{dx}$$

Generalised Derivative of Logarithmic Functions

not given
in exam

$\Rightarrow$

$$\frac{d}{dx}(\log_b u) = \frac{1}{(\ln b)u} \cdot \frac{du}{dx}$$

proof $\frac{d}{dx}(\log_b u) = \frac{d}{dx} \left[\frac{\ln u}{\ln b} \right] = \frac{1}{\ln b} \frac{d}{dx} \ln u$
 $= \frac{1}{\ln b} \cdot \frac{1}{u} \frac{du}{dx}$

$\therefore$ This expression is useful but if you don't remember, you can derive it!!

Ex. 4: Show that the derivative of $f(x) = x - \ln \sqrt{x}$ is $f'(x) = 1 - \frac{1}{2x}$. In addition, show

that an equation for the tangent line to the graph of $f(x) = x - \ln \sqrt{x}$ at the point where $x = 1$ is $y = 0.5x + 0.5$.

Derivative of Exponential Functions to the Base 'a' (a is a real number)

We can use the formula for the derivatives of e^u to differentiate a more general exponential function a^u . Replacing a with its equivalent from $e^{\ln a}$. $[a = e^{\ln a}]$

$$\begin{aligned} \frac{d(a^u)}{dx} &= \frac{d}{dx} \left[(e^{\ln a})^u \right] \\ &= \frac{d}{dx} (e^{u \ln a}) \\ &= e^{u \ln a} \frac{d}{dx} (u \ln a) \\ &= e^{u \ln a} \left(\frac{du}{dx} \right) (\ln a) \end{aligned}$$

This substitution is useful in differentiation of a function in power form
 e.g. $y = a^u$
 where a is a constant

not given
in exam

$\Rightarrow$

$$\frac{d(a^u)}{dx} = a^u \ln a \left(\frac{du}{dx} \right)$$

both of these expressions will not be given⁵ in exam so I do not expect students to remember them. What I expect you to understand is that if you have to diff a log function, you will have to try to change it into $\ln$ form.

Similarly, when you have to diff a power function a^u , you will have to change $a = e^{\ln a}$

This example will help you understand how to diff a power function by substitute

Ex. 4: Determine the derivative of the following functions.

(a) $y = a^x$.

Ans: $y' = a^x \ln a$

(b) $y = 20^x$

Ans: $y' = 20^x \ln 20$

(c) $y = x2^{3x}$

Ans: $y' = 2^{3x}(1 + 3x \ln 2)$

(d) $y = \alpha 10^{-x} + \beta 10^{-x^2}$

Ans: $y' = -\ln 10(\alpha 10^{-x} + 2\beta x 10^{-x^2})$

$a = e^{\ln a}$

where a is constant

(complete solution in separate file)

5.4 Logarithmic Differentiation

Technique of substitute $a = e^{\ln a}$

This technique is often useful to simplify the differentiation of $y = f(x)$ when $f(x)$ involves products, quotients or powers.

Ex. 5: Show that the derivative of $y = 2^x$ is $y' = (\ln 2)2^x$.

Ex. 6: Show that the derivative of $y = (x+1)^2(x+2)$ is $y' = (x+1)(3x+5)$.

Ex. 7: Show that the derivative of $y = x^{2x+1}$ is $y' = x^{2x+1} \left[2 \ln x + \frac{2x+1}{x} \right]$.


Try Ex 5, 6, 7, you'll see that substitute $a = e^{\ln a}$ is most useful when you have to diff a function in power form where the power contains variables such as x in Ex 5 and $2x+1$ in Ex. 7

5.5 Curve Sketching

Not in exam but give Ex. 8 as a homework!! (given marks)

Ex. 8: $f(x) = x^2 - 8 \ln x$ intercepts the x axis at $x \approx 1.2$ and $x \approx 2.9$. Sketch the graph of

$f(x) = x^2 - 8 \ln x$.



Ans: Domain is positive real number.

$$f'(x) = \frac{2x^2 - 8}{x}$$


$f(x)$ is decreasing between $(0, 2)$ and is increasing between $(2, \infty)$. The minimum point is $(2, -1.5)$. The function is always concave up and there is no inflection

point. $x = 0$ is the vertical asymptote and there is no horizontal asymptote.

Ex. 9: Determine where the function

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$

is increasing and decreasing, and where its graph is concave up and concave down. Find the relative extrema and inflection points and draw the graph.



Ans: $f'(x) = \frac{-x}{\sqrt{2\pi}} e^{-\frac{x^2}{2}},$

$f''(x) = \frac{-x}{\sqrt{2\pi}} (x^2 - 1) e^{-\frac{x^2}{2}}$ (0, 0.4) is the relative maximum. (1, 0.24) and (-1, 0.24) are inflection points. The function is increasing for negative x and decreasing for positive x . The function is concave up where $x < -1$ and $x > 1$; and concave down where $-1 < x < 1$. There is no x intercept. y

$= 0$ is the horizontal asymptote.