

PROBLEMS

C.1 Let $Y_1, Y_2, Y_3,$ and Y_4 be independent, identically distributed random variables from a population with mean μ and variance σ^2 . Let $\bar{Y} = \frac{1}{4}(Y_1 + Y_2 + Y_3 + Y_4)$ denote the average of these four random variables.

- (i) What are the expected value and variance of \bar{Y} in terms of μ and σ^2 ?
- (ii) Now, consider a different estimator of μ :

$$W = \frac{1}{8}Y_1 + \frac{1}{8}Y_2 + \frac{1}{4}Y_3 + \frac{1}{2}Y_4.$$

This is an example of a *weighted* average of the Y_i . Show that W is also an unbiased estimator of μ . Find the variance of W .

- (iii) Based on your answers to parts (i) and (ii), which estimator of μ do you prefer, \bar{Y} or W ?

C.2 This is a more general version of Problem C.1. Let Y_1, Y_2, \dots, Y_n be n pairwise uncorrelated random variables with common mean μ and common variance σ^2 . Let \bar{Y} denote the sample average.

- (i) Define the class of *linear estimators* of μ by

$$W_a = a_1Y_1 + a_2Y_2 + \dots + a_nY_n,$$

where the a_i are constants. What restriction on the a_i is needed for W_a to be an unbiased estimator of μ ?

- (ii) Find $\text{Var}(W_a)$.
- (iii) For any numbers a_1, a_2, \dots, a_n , the following inequality holds: $(a_1 + a_2 + \dots + a_n)^2/n \leq a_1^2 + a_2^2 + \dots + a_n^2$. Use this, along with parts (i) and (ii), to show that $\text{Var}(W_a) \geq \text{Var}(\bar{Y})$ whenever W_a is unbiased, so that \bar{Y} is the *best linear unbiased estimator*. [Hint: What does the inequality become when the a_i satisfy the restriction from part (i)?]

C.3 Let \bar{Y} denote the sample average from a random sample with mean μ and variance σ^2 . Consider two alternative estimators of μ : $W_1 = [(n-1)/n]\bar{Y}$ and $W_2 = \bar{Y}/2$.

- (i) Show that W_1 and W_2 are both biased estimators of μ and find the biases. What happens to the biases as $n \rightarrow \infty$? Comment on any important differences in bias for the two estimators as the sample size gets large.
- (ii) Find the probability limits of W_1 and W_2 . [Hint: Use Properties PLIM.1 and PLIM.2; for W_1 , note that $\text{plim} [(n-1)/n] = 1$.] Which estimator is consistent?
- (iii) Find $\text{Var}(W_1)$ and $\text{Var}(W_2)$.
- (iv) Argue that W_1 is a better estimator than \bar{Y} if μ is "close" to zero. (Consider both bias and variance.)

C.4 For positive random variables X and Y , suppose the expected value of Y given X is $E(Y|X) = \theta X$. The unknown parameter θ shows how the expected value of Y changes with X .

- (i) Define the random variable $Z = Y/X$. Show that $E(Z) = \theta$. [Hint: Use Property CE.2 along with the law of iterated expectations, Property CE.4. In particular, first show that $E(Z|X) = \theta$ and then use CE.4.]
- (ii) Use part (i) to prove that the estimator $W_1 = n^{-1} \sum_{i=1}^n (Y_i/X_i)$ is unbiased for θ , where $\{(X_i, Y_i): i = 1, 2, \dots, n\}$ is a random sample.