
Instructions

6304641191 Pornchanok Thawornphiboon

- (1) Please read the instruction carefully. Also take this habit with you into the exam room.
- (2) Please read each question carefully and answer the questions straightforwardly. Always provide economic reasons at least a paragraph for your analysis, or a graph when necessary, even when the question does not indicate so.
- (3) Handing and submitting assignments are only available via BE Moodle.

Answering the questions and preparing answer sheets

- (1) Answers are to be handwritten, in either digital or analog form, in a blank canvas or any clean paper. Make sure that your handwriting is clearly visible and readable.
- (2) There is no need to rewrite the question. Just indicate the question number clearly for each of the answer, such as 1.a).
- (3) Default decimal point is 4.
- (4) Choose precise wordings, especially when you want to interpret the meaning of a test, confidence interval, or coefficients.
- (5) When done, for the digital case, collage all the pages into a single PDF file. For those who write on sheets of paper, take photo of all pages then convert all of them into a single PDF file as well.
- (6) Name your PDF file as StudentID_YourNickname, such as 640123456_Bo.

Submitting your answers

- (1) Make sure your file does not exceed 10MB. This is the maximum file size for BE Moodle upload.
- (2) Login to BE Moodle, head into the course, then the assignment topic.
- (3) Choose your file to submit. Done. There will be timestamp for your upload date and time, so please make sure to not submit later than that.

1. (15 points) Given this information,

$n = 46$	$\sum_{i=1}^n X_i = 3,959.80$	$\sum_{i=1}^n Y_i = 3,180.80$
$\bar{X} = 86.0826$	$\bar{Y} = 69.1478$	
$\sum_{i=1}^n (X_i)^2 = 364,023.30$		$\sum_{i=1}^n X_i Y_i = 319,943.18$
$\sum_{i=1}^n (X_i - \bar{X})^2 = 23,153.3861$		$\sum_{i=1}^n (Y_i - \bar{Y})^2 = 94,525.1748$
$\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y}) = 46,131.6183$		$\sum_{i=1}^n \hat{u}_i^2 = 2,610.9211$

answer the following questions. Show your work.

- (4 points) From regression model: $Y_i = \beta_1 + \beta_2 X_i + u_i$, $u_i \sim NIID(0, \sigma^2)$, find the estimators of β_1 and β_2 with OLS method and explain the meaning of the model.
- (2 points) Find R^2 and explain its meaning.
- (1 points) If $X_i = 60$, estimate the value of \hat{Y}_i and explain its meaning.
- (3 points) Find the estimators of $\text{var}(u_i)$, $\text{var}(\hat{\beta}_1)$ and $\text{var}(\hat{\beta}_2)$
- (2.5 points) What are the 95-percent confident intervals for β_2 ? Interpret the meaning.
- (2.5 points) Test the hypothesis whether coefficients (both β_1 and β_2) are different from zero at 0.05 level of significance.

Assignment 1

Assigned on Sep 14th, 2021. Due date Sep 27th, 2021 before midnight.

2. (8 points) Answer the following question without any mathematical proof. A 3-6-line paragraph for a question is sufficient.
- (2 points) If we have only one data point, can we create a sample regression function? Why?
 - (2 points) Does a significant β_2 sufficient for us to believe that X and Y are causally related? Provide an example to support your answer.
 - (2 points) When we test a hypothesis and find that β_2 is significantly different from zero, what does the result actually suggest? Answer with specific wording from statistical perspective.
 - (2 points) What is (are) (an) advantage(s) of an interval estimation over point estimate?
3. (7 points) Given that the dependent variable is natural log of wage (lwage) in Thai Baht and the independent variable is hours worked per week (main_hr), the result of estimation is shown in the table below here.

Source	SS	df	MS	Number of obs	=	308
Model	50.060869	1	50.060869	F(1, 306)	=	92.20
Residual	166.152715	306	.542982728	Prob > F	=	0.0000
				R-squared	=	0.2315
				Adj R-squared	=	0.2290
Total	216.213584	307	.704278775	Root MSE	=	.73687

lwage	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
main_hr	.0318017	.003312	9.60	0.000	.0252844 .0383189
_cons	7.658082	.1256392	60.95	0.000	7.410856 7.905308

Answer the following questions. Show your work.

- (2 points) On average, how much is the nominal wage for a person who works 0 hour a week? (Note that this is a point estimation, not a prediction)
- (2 points) If a person works an hour more, how much, on average, wage change do we expect?
- (3 points) If you want to change the reading of unit from hours worked to days worked, what values in the main_hr row will differ? Calculate the changes to all the values in that row, **disregarding the constant row**. You might want to impose an assumption here. State that clearly before calculation.

1. (15 points) Given this information,

$n = 46$	$\sum_{i=1}^n X_i = 3,959.80$	$\sum_{i=1}^n Y_i = 3,180.80$
$\bar{X} = 86.0826$	$\bar{Y} = 69.1478$	
$\sum_{i=1}^n (X_i)^2 = 364,023.30$	$\sum_{i=1}^n X_i Y_i = 319,943.18$	
$\sum_{i=1}^n (X_i - \bar{X})^2 = 23,153.3861$	$\sum_{i=1}^n (Y_i - \bar{Y})^2 = 94,525.1748$	
$\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y}) = 46,131.6183$	$\sum_{i=1}^n \hat{u}_i^2 = 2,610.9211$	

$\downarrow \sum (Y_i - \hat{Y}_i)^2$

answer the following questions. Show your work.

a) (4 points) From regression model: $Y_i = \beta_1 + \beta_2 X_i + u_i$, $u_i \sim NIID(0, \sigma^2)$, find the estimators of β_1 and β_2 with OLS method and explain the meaning of the model.

$$\hat{\beta}_2 = \frac{\sum (X_i - \bar{X})(Y_i - \bar{Y})}{\sum (X_i - \bar{X})^2} = \frac{46,131.6183}{23,153.3861} = 1.9924 \#$$

$$\hat{\beta}_1 = \bar{Y} - \hat{\beta}_2 \bar{X} = 69.1478 - (1.9924 \times 86.0826) = 69.1478 - 171.5109 = -102.3631 \#$$

Plugging the estimators into the equation, we have

$$\hat{Y} = \hat{\beta}_1 + \hat{\beta}_2 X_i$$

$$\hat{Y}_i = -102.3631 + 1.9924 X_i \quad \text{SRF}$$

The intercept of this model is at -102.361 and the slope is 1.9924.

When X_i increases by 1 unit, \hat{Y}_i increases by 1.9924. #

b) (2 points) Find R^2 and explain its meaning.

Note: $r^2 =$ coefficient of determination

$$R^2 = 1 - \frac{\sum \hat{u}_i^2}{\sum (Y_i - \bar{Y})^2} = 1 - \frac{2,610.9211}{94,525.1748} = 1 - 0.0276 = 0.9724 = 97.24\%$$

$1 - (RSS/TSS)$ Fits well b/c r^2 approaches $\rightarrow 1$.

R^2 Values of 0.9724 suggests that X_i explain about 97.24% of the variation in Y_i .

c) (1 points) If $X_i = 60$, estimate the value of \hat{Y}_i and explain its meaning.

Replace $X_i = 60$ to find \hat{Y}_i

$$\hat{Y}_i = -102.3631 + 1.9924(60) = 17.1809$$

When $X_i = 60$, \hat{Y}_i is 17.1809.

d) (3 points) Find the estimators of $\text{var}(u_i)$, $\text{var}(\hat{\beta}_1)$ and $\text{var}(\hat{\beta}_2)$

$$\text{var}(u_i) = \frac{\sum \hat{u}_i^2}{n-k} = \frac{\sum (Y_i - \hat{Y})^2}{n-k} \quad ; \quad \begin{array}{l} n \text{ is \# of observation} \\ k \text{ is \# of parameters estimated. } (\beta_1, \beta_2) \end{array}$$

$$= \frac{2610.9211}{46-2}$$



$$= 59.3391 \#$$

$$\text{var}(\hat{\beta}_1) = \frac{\sum X_i^2}{n \sum (X_i - \bar{X})^2} (\text{var}(u_i)) = \frac{364,023.30}{46(23153.3861)} \times 59.3391 = 20.2874 \#$$

$$\text{var}(\hat{\beta}_2) = \frac{\text{var}(u_i)}{\sum (X_i - \bar{X})^2} = \frac{59.3391}{23,153.3861} = 0.0026 \#$$

e) (2.5 points) What are the 95-percent confident intervals for β_2 ? Interpret the meaning.

$\alpha = 0.05$ we have only 1 estimator, we will use t-stat instead of z.



$$t = \frac{\hat{\beta}_2 - \beta_2}{\sigma_{\hat{\beta}_2}}$$

Degree of freedom = $46 - 2 = 44$

$$P\left[-t_{\alpha/2} < \frac{\hat{\beta}_2 - \beta_2}{\sigma_{\hat{\beta}_2}} < t_{\alpha/2}\right] = 1 - \alpha \rightarrow 0.05$$

$$P\left[\hat{\beta}_2 - (t_{\alpha/2})(\sigma_{\hat{\beta}_2}) < \beta_2 < \hat{\beta}_2 + (t_{\alpha/2})(\sigma_{\hat{\beta}_2})\right] = 95\%$$

$$P\left[(1.9924) - (2.021)(4.5035) < \beta_2 < (1.9924) + (2.021)(4.5035)\right] = 95\%$$

$$P\left[1.9924 - 9.1016 < \beta_2 < 1.9924 + 9.1016\right] = 95\%$$

$$P\left[-7.1092 < \beta_2 < 11.094\right] = 95\%$$

\therefore We can interpret that β_2 can be between -7.1092 to 11.094 which 95% of observation β_2 fall in this interval.

f) (2.5 points) Test the hypothesis whether coefficients (both β_1 and β_2) are different from zero at 0.05 level of significance.

2. (8 points) Answer the following question without any mathematical proof. A 3-6-line paragraph for a question is sufficient.

- a) (2 points) If we have only one data point, can we create a sample regression function? Why?
- b) (2 points) Does a significant β_2 sufficient for us to believe that X and Y are causally related? Provide an example to support your answer.
- c) (2 points) When we test a hypothesis and find that β_2 is significantly different from zero, what does the result actually suggest? Answer with specific wording from statistical perspective.
- d) (2 points) What is (are) (an) advantage(s) of an interval estimation over point estimate?

a) No, if we have only one data point, the SRF can be many slope and SRF is a linear line and a linear line need data points at least 2 points.

b) A significant β_2 can tell us How X and Y are causally related because β_2 is slope of PRF
 If $\beta_2 = 0$ or no β_2 , PRF will be horizontal line.
 $\beta_2 \neq 0$, when x increase by 1, dependent variable or Y increase by a .
 or a

c) When β_2 is not zero, we can make sure that X is related to Y .

d) When we try to find the estimators, if it came up in point which is single number.

There will be error because when we draw the sample, the group of data can be vary. Therefore, if we estimate them in interval, the data interval can cover the range of data that can occur.

3. (7 points) Given that the dependent variable is natural log of wage (lwage) in Thai Baht and the independent variable is hours worked per week (main_hr), the result of estimation is shown in the table below here.

Source	SS	df	MS	Number of obs	=	308
ESS				F(1, 306)	=	92.20
RSS	Model	1	50.060869	Prob > F	=	0.0000
TSS	Residual	306	.542982728	R-squared	=	0.2315
	Total	307	.704278775	Adj R-squared	=	0.2290
				Root MSE	=	.73687

	lwage	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
indep	main_hr	.0318017	.003312	9.60	0.000	.0252844 .0383189
depond	_cons	7.658082	.1256392	60.95	0.000	7.410856 7.905308

Answer the following questions. Show your work.

- a) (2 points) On average, how much is the nominal wage for a person who works 0 hour a week? (Note that this is a point estimation, not a prediction)
- b) (2 points) If a person works an hour more, how much, on average, wage change do we expect?
- c) (3 points) If you want to change the reading of unit from hours worked to days worked, what values in the main_hr row will differ? Calculate the changes to all the values in that row, **disregarding the constant row**. You might want to impose an assumption here. State that clearly before calculation.

$$\ln \widehat{wage} = \beta_1 + \beta_2(\text{main_hr})$$

$$a) \hat{\beta}_1 = 7.658082$$

b) Wage will increase by $100(0.0318017)$
 or 3.18017%

c) hr ↓ day
 $\ln \widehat{wage} = 7.658082 + 0.0318017(\text{main_hour})$; If work 0 hr, get 7.658082 THB
 $se = (0.1256392) (0.00318017)$; work 1 more hr, get 3.18017 %
 $\ln \widehat{wage} = 7.658082 + (0.0318017)(24)(\text{main_day})$; if work 0 day, get 7.658082 THB
 $se = (0.1256392) (0.003312)(24)$; work 1 more day, get 36.3240 %
 $\sigma_{\hat{\beta}_1}$ $\sigma_{\hat{\beta}_2}$