

## Matrix inverses

The inverse of a real number  $a$  is denoted by  $a^{-1}$ . For example,  $7^{-1} = 1/7$  and

$$7 \cdot 7^{-1} = 7^{-1} \cdot 7 = 1$$

An  $n \times n$  matrix  $A$  is said to be **invertible** if there is an  $n \times n$  matrix  $C$  satisfying

$$CA = AC = I_n$$

where  $I_n$  is the  $n \times n$  identity matrix. We call  $C$  the **inverse** of  $A$ .

example  $A = \begin{bmatrix} 2 & 5 \\ -3 & -7 \end{bmatrix}$  ;  $C = \begin{bmatrix} -7 & -5 \\ 3 & 2 \end{bmatrix}$   $AC =$   $CA =$

The inverse of  $A$  is usually denoted by  $A^{-1}$ .

We have

$$AA^{-1} = A^{-1}A = I_n$$

**Not all  $n \times n$  matrices are invertible.** A matrix which is not invertible is sometimes called a **singular** matrix. An invertible matrix is called **nonsingular** matrix.

Dr. Julaluk Carmai

**Fact 1** If  $A$  is invertible, then the inverse is unique.

*Proof:* Assume  $B$  and  $C$  are both inverses of  $A$ . Then

$$B = BI = B(\text{---}) = (\text{---})\text{---} = I\text{---} = C.$$

So the inverse is unique since any two inverses coincide. ■

**Fact 2** The inverse of  $A^{-1}$  is  $A$  itself.

**Fact 3** Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . If  $ad - bc \neq 0$ , then  $A$  is invertible and

$$A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

If  $ad - bc = 0$ , then  $A$  is not invertible.

Dr. Julaluk Carmai

Assume  $A$  is any invertible matrix and we wish to solve  $AX = \mathbf{b}$ . Then

$$\text{---}AX = \text{---}\mathbf{b} \quad \text{and so}$$

$$IX = \text{---} \quad \text{or} \quad X = \text{---}.$$

Suppose  $\mathbf{w}$  is also a solution to  $AX = \mathbf{b}$ . Then  $A\mathbf{w} = \mathbf{b}$  and

$$\text{---}A\mathbf{w} = \text{---}\mathbf{b} \quad \text{which means} \quad \mathbf{w} = A^{-1}\mathbf{b}.$$

So,  $\mathbf{w} = A^{-1}\mathbf{b}$ , which is in fact the same solution.

We have proved the following result:

**Fact 4** If  $A$  is an invertible  $n \times n$  matrix, then for each  $\mathbf{b}$  in  $\mathbf{R}^n$ , the equation  $AX = \mathbf{b}$  has the unique solution  $\mathbf{x} = A^{-1}\mathbf{b}$ .

Dr. Julaluk Carmai

**EXAMPLE:** Use the inverse of  $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$  to solve

$$\begin{cases} -7x_1 + 3x_2 = 2 \\ 5x_1 - 2x_2 = 1 \end{cases}.$$

*Solution:* Matrix form of the linear system:

$$\begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{14-15} \begin{bmatrix} -2 & -3 \\ -5 & -7 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix}.$$

$$\mathbf{x} = A^{-1}\mathbf{b} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 14 \\ 19 \end{bmatrix}$$

Dr. Julaluk Carmai

### Properties of Inverses

Suppose A and B are invertible. Then the following results hold:

- a.  $A^{-1}$  is invertible and  $(A^{-1})^{-1} = A$  (i.e.  $A$  is the inverse of  $A^{-1}$ ).
- b.  $AB$  is invertible and  $(AB)^{-1} = B^{-1}A^{-1}$
- c.  $A^T$  is invertible and  $(A^T)^{-1} = (A^{-1})^T$

Partial proof of part b:

$$\begin{aligned} (AB)(B^{-1}A^{-1}) &= A(\text{_____})A^{-1} \\ &= A(\text{_____})A^{-1} = \text{_____} = \text{_____}. \end{aligned}$$

Similarly, one can show that  $(B^{-1}A^{-1})(AB) = I$ .

Proof part c

$$\begin{aligned} A^T(A^{-1})^T &= (A^{-1}A)^T = I^T = I \\ (A^{-1})^T A^T &= (AA^{-1})^T = I^T = I \end{aligned}$$

### Matrix inversion algorithm

$$AA^{-1} = I$$

$$\begin{bmatrix} 1 & 3 \\ 2 & 7 \end{bmatrix} \begin{bmatrix} a & c \\ b & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

→ Taken a column at a time, that equation determines the columns of  $A^{-1}$

$A$  times column  $j$  of  $A^{-1}$  = column  $j$  of  $I$

$$\begin{bmatrix} 1 & 3 \\ 2 & 7 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad ; \quad \begin{bmatrix} 1 & 3 \\ 2 & 7 \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$\left[ \begin{array}{cc|cc} 1 & 3 & 1 & 0 \\ 2 & 7 & 0 & 1 \end{array} \right]$$

Carry out elimination on all systems simultaneously.

$$\begin{aligned} &\left[ \begin{array}{cc|cc} 1 & 3 & 1 & 0 \\ 2 & 7 & 0 & 1 \end{array} \right] \\ &\downarrow \\ &\left[ \begin{array}{cc|cc} 1 & 3 & 1 & 0 \\ 0 & 1 & -2 & 1 \end{array} \right] \\ &\downarrow \\ &\left[ \begin{array}{cc|cc} 1 & 0 & 7 & -3 \\ 0 & 1 & -2 & 1 \end{array} \right] \\ &\downarrow \\ &\left[ \begin{array}{cc|cc} I & & & A^{-1} \end{array} \right] \end{aligned}$$

### The Gauss-Jordan method

### Matrix inversion algorithm

Place  $A$  and  $I$  side-by-side to form an augmented matrix  $[A \ I]$ . Then perform row operations on this matrix (which will produce identical operations on  $A$  and  $I$ ). So by Theorem 7:

$$[A \ I] \text{ will row reduce to } [I \ A^{-1}]$$

or  $A$  is not invertible.

**EXAMPLE:** Find the inverse of  $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ , if it exists.

Solution:

$$[A \ I] = \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \sim \dots \sim \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right]$$

$$\text{So } A^{-1} = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & 0 & 1 \\ \frac{3}{2} & 1 & 0 \end{bmatrix}$$

Find the inverse of the matrix

$$A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 0 & 3 \\ 4 & -3 & 8 \end{bmatrix}, \text{ if it exists}$$

Dr. Julaluk Carmai

## Elementary Matrices

### Definition

An **elementary matrix** is one that is obtained by performing a single elementary row operation on an identity matrix.

**EXAMPLE:** Let  $E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ,  $E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ ,

$$E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} \text{ and } A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}.$$

Dr. Julaluk Carmai

Observe the following products and describe how these products can be obtained by elementary row operations on  $A$ .

$$E_1 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ 2d & 2e & 2f \\ g & h & i \end{bmatrix}$$

$$E_2 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ g & h & i \\ d & e & f \end{bmatrix}$$

$$E_3 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 3a+g & 3b+h & 3c+i \end{bmatrix}$$

If an elementary row operation is performed on an  $m \times n$  matrix  $A$ , the resulting matrix can be written as  $EA$ , where the  $m \times m$  matrix  $E$  is created by performing the same row operations on  $I_m$ .

Dr. Julaluk Carmai

Elementary matrices are *invertible* because row operations are *reversible*. To determine the inverse of an elementary matrix  $E$ , determine the elementary row operation needed to transform  $E$  back into  $I$  and apply this operation to  $I$  to find the inverse.

For example,

$$E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} \quad E_3^{-1} = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$$

Dr. Julaluk Carmai

**Example:** Let  $A = \begin{bmatrix} 1 & 0 & 0 \\ -\frac{3}{2} & 0 & \frac{1}{2} \\ 0 & 1 & 0 \end{bmatrix}$ . Then

$$E_1 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -\frac{3}{2} & 0 & \frac{1}{2} \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

$$E_2(E_1 A) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix}$$

$$E_3(E_2 E_1 A) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

So

$$E_3 E_2 E_1 A = I_3.$$

Then multiplying on the right by  $A^{-1}$ , we get

$$E_3 E_2 E_1 A \underline{\hspace{1cm}} = I_3 \underline{\hspace{1cm}}.$$

So

$$E_3 E_2 E_1 I_3 = A^{-1}$$

**The elementary row operations that row reduce  $A$  to  $I_n$  are the same elementary row operations that transform  $I_n$  into  $A^{-1}$ .**

### Matrix Factorisations

A **factorisation** of a matrix  $A$  is an equations that expresses  $A$  as a product of 2 or more matrices.

Gauss elimination can be used to find such factorisations.

$$Ax=b \quad \rightarrow \quad Ux=c \quad Ax = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 5 \\ -2 \\ 9 \end{bmatrix}$$

How to relate  $A$  to  $U$ ?

E.g.

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix} \quad \rightarrow \quad U = \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

### 3 elimination steps

- i) Subtract 2 times the first equation from the second;
- ii) Subtract  $-1$  times the first equation from the third;
- iii) Subtract  $-1$  times the second equation from the third

Each step corresponds to an elementary matrix.

Let matrix  $E$  for step i),  $F$  for step ii) and  $G$  for step iii)

$$E = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad ; \quad F = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \quad ; \quad G = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$

$$GFEA = U$$

The single matrix that take  $A$  to  $U$  is  $GFE$

$$GFE = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix}$$

$$GFEA = U$$

$$A = (GFE)^{-1}U = E^{-1}F^{-1}G^{-1}U$$

$$E^{-1}F^{-1}G^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & -1 & 1 \end{bmatrix} = L \quad \text{L= lower triangular}$$

$$A = LU \rightarrow \text{Triangular (LU) factorisation}$$

The entries below the diagonal are exactly the multipliers  $l_{21}=2, -1$  and  $l_{31}=-1$ .

If no row exchanges are required, the original matrix  $A$  can be written as a product  $A=LU$ . The matrix  $L$  is lower triangular, with 1's on the diagonal and the multipliers  $l_{ij}$  (taken from elimination ) below the diagonal.  $U$  is the upper triangular matrix which appears after forward elimination and before back substitution; its diagonal entries are pivots.

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \rightarrow U = \begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix} \quad \text{with} \quad L = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 0 & 2 \\ 3 & 4 \end{bmatrix} \rightarrow \text{Needs a row exchange} \\ \text{cannot be factored into } A=LU$$

**Remark** You have to be careful with  $L$ . Suppose elimination subtracts row 1 from row 2, creating  $l_{21}$ . Then suppose it exchanges rows 2 and 3. If that exchange is done in advance, the multiplier will change to  $l_{31}$  in  $PA=LU$

$$\text{E.g. } A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 3 \\ 2 & 5 & 8 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 0 & 0 & 2 \\ 0 & 3 & 6 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 0 & 3 & 6 \\ 0 & 0 & 2 \end{bmatrix} = U$$

With rows exchanged, we recover LU –but now  $l_{31}=1$  and  $l_{21}=2$ :

$$P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad \text{and} \quad L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

$$A_{m \times n} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ * & 1 & 0 & 0 \\ * & * & 1 & 0 \\ * & * & * & 1 \end{bmatrix} \begin{bmatrix} \bullet & * & * & * & * \\ 0 & \bullet & * & * & * \\ 0 & 0 & 0 & \bullet & * \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \\ \begin{matrix} L \\ m \times m \end{matrix} \quad \begin{matrix} U \\ m \times n \end{matrix}$$

**EXAMPLE 1** It can be verified that

$$A = \begin{bmatrix} 3 & -7 & -2 & 2 \\ -3 & 5 & 1 & 0 \\ 6 & -4 & 0 & -5 \\ -9 & 5 & -5 & 12 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 2 & -5 & 1 & 0 \\ -3 & 8 & 3 & 1 \end{bmatrix} \begin{bmatrix} 3 & -7 & -2 & 2 \\ 0 & -2 & -1 & 2 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & -1 \end{bmatrix} = LU$$

Use this LU factorization of  $A$  to solve  $Ax = b$ , where  $b = \begin{bmatrix} -9 \\ 5 \\ 7 \\ 11 \end{bmatrix}$ .

$$[L \quad \mathbf{b}] = \begin{bmatrix} 1 & 0 & 0 & 0 & -9 \\ -1 & 1 & 0 & 0 & 5 \\ 2 & -5 & 1 & 0 & 7 \\ -3 & 8 & 3 & 1 & 11 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 0 & -9 \\ 0 & 1 & 0 & 0 & -4 \\ 0 & 0 & 1 & 0 & 5 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix} = [I \quad \mathbf{y}]$$

$$[U \quad \mathbf{y}] = \begin{bmatrix} 3 & -7 & -2 & 2 & -9 \\ 0 & -2 & -1 & 2 & -4 \\ 0 & 0 & -1 & 1 & 5 \\ 0 & 0 & 0 & -1 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 0 & 3 \\ 0 & 1 & 0 & 0 & 4 \\ 0 & 0 & 1 & 0 & -6 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} 3 \\ 4 \\ -6 \\ -1 \end{bmatrix}$$

### Algorithm for an LU factorization

1. Reduce  $\mathbf{A}$  to an echelon form  $\mathbf{U}$  by a sequence of row replacement operations, if possible.
2. Place entries in  $\mathbf{L}$  such that the same sequence of row operations reduces  $\mathbf{L}$  to  $\mathbf{I}$

Dr. Julaluk Carmai

EXAMPLE 2 Find an LU factorization of

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ -4 & -5 & 3 & -8 & 1 \\ 2 & -5 & -4 & 1 & 8 \\ -6 & 0 & 7 & -3 & 1 \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ -4 & -5 & 3 & -8 & 1 \\ 2 & -5 & -4 & 1 & 8 \\ -6 & 0 & 7 & -3 & 1 \end{bmatrix} \sim \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ 0 & -3 & 1 & 2 & -3 \\ 0 & -9 & -3 & -4 & 10 \\ 0 & 12 & 4 & 12 & -5 \end{bmatrix} = \mathbf{A}_1$$

$$\sim \mathbf{A}_2 = \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ 0 & 3 & 1 & 2 & -3 \\ 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 4 & 7 \end{bmatrix} \sim \begin{bmatrix} 2 & 4 & -1 & 5 & -2 \\ 0 & 3 & 1 & 2 & -3 \\ 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 5 \end{bmatrix} = \mathbf{U}$$

$$\mathbf{L} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 1 & & 1 & 0 \\ -3 & & & 1 \end{bmatrix}$$

Dr. Julaluk Carmai

$$\begin{bmatrix} 2 \\ -4 \\ 2 \\ -6 \end{bmatrix} \begin{bmatrix} 3 \\ -9 \\ 12 \\ 4 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \end{bmatrix} \begin{bmatrix} 5 \\ 5 \end{bmatrix} \begin{matrix} \div 2 \\ \div 3 \\ \div 2 \\ \div 5 \end{matrix} \begin{matrix} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{matrix} \begin{bmatrix} 1 & & & \\ -2 & 1 & & \\ 1 & -3 & 1 & \\ -3 & 4 & 2 & 1 \end{bmatrix}, \quad \text{and} \quad \mathbf{L} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 1 & -3 & 1 & 0 \\ -3 & 4 & 2 & 1 \end{bmatrix}$$

An easy calculation verifies that this  $\mathbf{L}$  and  $\mathbf{U}$  satisfy  $\mathbf{LU} = \mathbf{A}$ .

Dr. Julaluk Carmai