

TWO-VARIABLE REGRESSION MODEL: THE PROBLEM OF ESTIMATION

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

The Two-Variable PRF:

$$Y_i = \beta_1 + \beta_2 X_i + u_i$$

The Two-Variable SRF:

$$Y_i = \hat{\beta}_1 + \hat{\beta}_2 X_i + \hat{u}_i$$

$$= \hat{Y}_i + \hat{u}_i$$

$\hat{Y}_i$ is the estimated (conditional mean) value of Y_i


ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

$$\hat{u}_i = Y_i - \hat{Y}_i$$

$$= Y_i - \hat{\beta}_1 - \hat{\beta}_2 X_i$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

ORDINARY LEAST SQUARES (OLS)



ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

ORDINARY LEAST SQUARES (OLS)

$$\sum \hat{u}_i^2 = \sum (Y_i - \hat{Y}_i)^2$$

$$= \sum (Y_i - \hat{\beta}_1 - \hat{\beta}_2 X_i)^2$$

The principle or the method of least squares chooses $\hat{\beta}_1$ and $\hat{\beta}_2$ in such a manner that, for a given sample or set of data, $\sum \hat{u}_i^2$ is as small as possible

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

$$\frac{\partial (\sum \hat{u}_i^2)}{\partial \hat{\beta}_1} = -2 \sum (Y_i - \hat{\beta}_1 - \hat{\beta}_2 X_i) = -2 \sum \hat{u}_i$$

$$\frac{\partial (\sum \hat{u}_i^2)}{\partial \hat{\beta}_2} = -2 \sum (Y_i - \hat{\beta}_1 - \hat{\beta}_2 X_i) X_i = -2 \sum \hat{u}_i X_i$$

Setting these equation to zero

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

$$\sum Y_i = n\hat{\beta}_1 + \hat{\beta}_2 \sum X_i$$

$$\sum Y_i X_i = \hat{\beta}_1 \sum X_i + \hat{\beta}_2 \sum X_i^2$$

Where n is the sample size. These simultaneous equations are known as the **Normal Equation**

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\begin{aligned}\hat{\beta}_2 &= \frac{n \sum X_i Y_i - \sum X_i \sum Y_i}{n \sum X_i^2 - (\sum X_i)^2} \\ &= \frac{\sum (X_i - \bar{X})(Y_i - \bar{Y})}{\sum (X_i - \bar{X})^2} \\ &= \frac{\sum x_i y_i}{\sum x_i^2}\end{aligned}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\begin{aligned}\hat{\beta}_2 &= \frac{\sum x_i y_i}{\sum x_i^2} \\ &= \frac{\sum x_i y_i}{\sum X_i^2 - n\bar{X}^2} \\ &= \frac{\sum X_i y_i}{\sum X_i^2 - n\bar{X}^2}\end{aligned}$$


ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\begin{aligned}\hat{\beta}_1 &= \frac{\sum X_i^2 \sum Y_i - \sum X_i \sum X_i Y_i}{n \sum X_i^2 - (\sum X_i)^2} \\ &= \bar{Y} - \hat{\beta}_2 \bar{X}\end{aligned}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

EXAMPLE ☺

Y	X
49	1
45	2
44	3
39	4
38	5
37	6
34	7
33	8
30	9



ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)


$$\begin{aligned}\hat{\beta}_2 &= \frac{\sum (X_i - \bar{X})(Y_i - \bar{Y})}{\sum (X_i - \bar{X})^2} = \frac{-178}{82.5} \approx -2.1576 \\ \hat{\beta}_1 &= \bar{Y} - \hat{\beta}_2 \bar{X} = 37.8 - (-2.1576)(5.5) = 49.667 \\ \bar{X} &= 5.5 \\ \bar{Y} &= 37.8\end{aligned}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

Y	X	$(X_i - \bar{X})$	$(X_i - \bar{X})^2$	$Y_i - \bar{Y}$	$(X_i - \bar{X})(Y_i - \bar{Y})$
49	1				
45	2				
44	3				
39	4				
38	5				
37	6				
34	7				
33	8				
30	9				
29	10				

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

$$\hat{Y}_i = 49.667 - 2.1576X_i$$



ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

PRACTICE I

A Random Sample from the Population
of Table 2.1

Weekly consumption expenditure \$ (Y)	Weekly income \$(X)
70	80
65	100
90	120
95	140
110	160
115	180
120	200
140	220
155	240
150	260

$$\hat{Y} = 0.5091X_i + 24.455$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

PRACTICE II

Another Random Sample from the
Population of Table 2.1


Weekly consumption expenditure \$ (Y)	Weekly income \$(X)
Y	X
55	80
88	100
90	120
80	140
118	160
120	180
145	200
135	220
145	240
175	260

$$\hat{Y} = 0.5761X_i + 17.17$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

THE REGRESSION LINE HAS THE FOLLOWING PROPERTIES:

- Passes through the sample means of Y and X



ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

- The mean value of the estimated $Y = \hat{Y}_i$ is equal to the mean value of the actual Y for

$$\hat{Y}_i = \hat{\beta}_1 + \hat{\beta}_2 X_i$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

CLASSICAL LINEAR REGRESSION MODEL (CLRM)

The assumptions underlying the method of least squares:

1. Linear regression model
2. Fixed X values or X values independent of the error term
3. Zero mean value of disturbance
4. Homoscedasticity or Constant Variance of u_i
5. No autocorrelation between the disturbances
6. The number of observations n must be greater than the number of parameters to be estimated
7. The nature of X variables

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

LINEAR REGRESSION MODEL

$$Y_i = \beta_1 + \beta_2 X_i + u_i$$

- Linear in the parameters
- May or may not be linear in the variables

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

X VALUES INDEPENDENT OF THE ERROR TERM


$$\text{Cov}(X_i, u_i) = 0$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

ZERO MEAN VALUE OF DISTURBANCE TERM

$$E(u_i | X_i) = 0$$

$$E(u_i) = 0$$




ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

HOMOSCEDASTICITY

$$\begin{aligned} \text{var}(u_i) &= E[u_i - E(u_i | X_i)]^2 \\ &= E(u_i^2 | X_i), \text{ because of assumption 3} \\ &= E(u_i^2), \text{ if } X_i \text{ are nonstochastic} \\ &= \sigma^2 \end{aligned}$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

HOMOSCEDASTICITY VS. HETEROSCEDASTICITY



HOMOSCEDASTICITY V.S. HETEROSCEDASTICITY

Homoscedasticity

- Equal variance
- The variation around the regression line is the same across the X values


Heteroscedasticity

- Unequal variance


ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

NO AUTOCORRELATION BETWEEN THE DISTURBANCES

Given X_i , the deviations of any two Y values from their mean value do not exhibit patterns such as those shown in figures below



NO AUTOCORRELATION BETWEEN THE DISTURBANCES



NO AUTOCORRELATION BETWEEN THE DISTURBANCES

Given any two X values, X_i and $X_j (i \neq j)$, the correlation between any two u_i and $u_j (i \neq j)$ is zero.

$$\text{cov}(u_i, u_j | X_i, X_j) = 0$$

$$\text{cov}(u_i, u_j) = 0, \text{ if } X \text{ is nonstochastic}$$

Where i and j are two different observation and where cov means covariance

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

THE NUMBER OF OBSERVATIONS N MUST BE GREATER THAN THE NUMBER OF PARAMETERS TO BE ESTIMATED

- The number of observations must be greater than the number of explanatory variables

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

THE NATURE OF X VARIABLES

- The X values in a given sample must not all be the same
- No outliers in the X values

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

STANDARD ERRORS OF LEAST-SQUARES ESTIMATES

$$\begin{aligned}\text{var}(\hat{\beta}_2) &= \frac{\sigma^2}{\sum x_i^2} \\ \text{se}(\hat{\beta}_2) &= \frac{\sigma}{\sqrt{\sum x_i^2}} \\ \text{var}(\hat{\beta}_1) &= \frac{\sum X_i^2}{n \sum x_i^2} \sigma^2 \\ \text{se}(\hat{\beta}_1) &= \sqrt{\frac{\sum X_i^2}{n \sum x_i^2}} \sigma\end{aligned}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\hat{\sigma}^2 = \frac{\sum \hat{u}_i^2}{n-2}$$

$\hat{\sigma}^2$ is the OLS estimator of the true but unknown σ^2

The expression n-2 is known as the number of degrees of freedom

$\sum \hat{u}_i^2$ is the sum of the residuals squared or residual sum of squares (RSS)

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\begin{aligned}\text{COV}(\hat{\beta}_1, \hat{\beta}_2) &= -\bar{X} \text{var}(\hat{\beta}_2) \\ &= -\bar{X} \left(\frac{\sigma^2}{\sum x_i^2} \right) \\ &= \frac{-\bar{X} \sigma^2}{\sum (X_i - \bar{X})^2}\end{aligned}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

EXAMPLE ☺

Y	X
49	1
45	2
44	3
39	4
38	5
37	6
34	7
33	8
30	9

$$\begin{aligned}\hat{\beta}_1 &= 49.667 \\ \hat{\beta}_2 &= -2.1576 \\ \bar{X} &= 5.5\end{aligned}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

Y	X	$\hat{Y}_i = \hat{\beta}_1 + \hat{\beta}_2 X_i$	$\hat{u}_i$	$\hat{u}_i^2$	$(X_i - \bar{X})^2$	X_i^2
49	1	47.5094	1.4906	2.2219	20.25	1
45	2	45.3518	-0.352	0.1238	12.25	4
44	3	43.1942	0.8058	0.6493	6.25	9
39	4	41.0366	-2.037	4.1477	2.25	16
38	5	38.879	-0.879	0.7726	0.25	25
37	6	36.7214	0.2786	0.0776	0.25	36
34	7	34.5638	-0.564	0.3179	2.25	49
33	8	32.4062	0.5938	0.3526	6.25	64
30	9	30.2486	-0.249	0.0618	12.25	81
29	10	28.091	0.909	0.8263	20.25	100
		Total		9.5515	82.5	385

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\hat{\sigma}^2 = \frac{\sum \hat{u}_i^2}{n-2} = \frac{9.5515}{10-2} = 1.1939$$

$$\text{var}(\hat{\beta}_2) = \frac{\hat{\sigma}^2}{\sum x_i^2} = \frac{\hat{\sigma}^2}{\sum (X_i - \bar{X})^2} = \frac{1.1939}{82.5} = 0.0145$$

$$\text{var}(\hat{\beta}_1) = \frac{\sum X_i^2}{n \sum x_i^2} \hat{\sigma}^2 = \frac{\sum X_i^2}{n \sum (X_i - \bar{X})^2} \hat{\sigma}^2 = \frac{(1.1939)(385)}{(10)(82.5)} = 0.5572$$

$$\text{COV}(\hat{\beta}_1, \hat{\beta}_2) = -\bar{X} \text{var}(\hat{\beta}_2) = -(5.5)(0.0145) = -0.07975$$


ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

PROPERTIES OF LEAST-SQUARES ESTIMATORS: THE GAUSS-MARKOV THEOREM

Best Linear Unbiased Estimator (BLUE) of β_2 :

- It is linear, that is, a linear function of a random variable, such as the dependent variable Y in the regression model
- It is unbiased $E(\hat{\beta}_2) = \beta_2$
- Efficient estimator – an unbiased estimator with the least variance


ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)



THE COEFFICIENT OF DETERMINATION r^2

- A measure of goodness of fit
- A summary measure that tells how well the sample regression line fits the data
- Measures the proportion or percentage of the total variation in Y explained by the regression model

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)




TO COMPUTE THIS r^2

$$Y_i = \hat{Y}_i + \hat{u}_i$$

$$y_i = \hat{y}_i + \hat{u}_i$$

$$\begin{aligned} \sum y_i^2 &= \sum \hat{y}_i^2 + \sum \hat{u}_i^2 + 2 \sum \hat{y}_i \hat{u}_i \\ &= \sum \hat{y}_i^2 + \sum \hat{u}_i^2 \\ &= \hat{\beta}_2^2 \sum x_i^2 + \sum \hat{u}_i^2 \end{aligned}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)



ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

$$\sum y_i^2 = \sum (Y_i - \bar{Y})^2 \quad \text{Total variation of the actual Y values about their sample mean}$$

(Total Sum of Squares, TSS)

$$\sum \hat{y}_i^2 = \sum (\hat{Y}_i - \bar{Y})^2 = \hat{\beta}_2^2 \sum x_i^2$$

Variation of the estimated Y values about their mean
(Explained Sum of Squares, ESS)

$$\sum \hat{u}_i^2 \quad \text{Residual or unexpected variation of the Y values about the regression line (Residual Sum of Squares, RSS)}$$

$$TSS = ESS + RSS$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

$$1 = \frac{ESS}{TSS} + \frac{RSS}{TSS}$$

$$= \frac{\sum (\hat{Y}_i - \bar{Y})^2}{\sum (Y_i - \bar{Y})^2} + \frac{\sum \hat{u}_i^2}{\sum (Y_i - \bar{Y})^2}$$

$$r^2 = \frac{\sum (\hat{Y}_i - \bar{Y})^2}{\sum (Y_i - \bar{Y})^2} = \frac{ESS}{TSS}$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

EXAMPLE ☺

Y	X
49	1
45	2
44	3
39	4
38	5
37	6
34	7
33	8
30	9

$$\hat{\beta}_1 = 49.667$$

$$\hat{\beta}_2 = -2.1576$$

$$\bar{X} = 5.5$$

$$\bar{Y} = 38$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

Y	X	$\hat{Y}_i = \hat{\beta}_1 + \hat{\beta}_2 X_i$	$\hat{u}_i$	$(Y_i - \bar{Y})^2$	$(\hat{Y}_i - \bar{Y})^2$	$\hat{u}_i^2$
49	1					
45	2					
44	3					
39	4					
38	5					
37	6					
34	7					
33	8					
30	9					
29	10					

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

Y	X	$\hat{Y}_i = \hat{\beta}_1 + \hat{\beta}_2 X_i$	$\hat{u}_i$	$(Y_i - \bar{Y})^2$	$(\hat{Y}_i - \bar{Y})^2$	$\hat{u}_i^2$
49	1	47.509				
45	2	45.352				
44	3	43.194				
39	4	41.037				
38	5	38.879				
37	6	36.721				
34	7	34.564				
33	8	32.406				
30	9	30.249				
29	10	28.091				

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

$$r^2 = \frac{\sum (\hat{Y}_i - \bar{Y})^2}{\sum (Y_i - \bar{Y})^2} = \frac{ESS}{TSS}$$

$$r^2 = \frac{\sum (\hat{Y}_i - \bar{Y})^2}{\sum (Y_i - \bar{Y})^2} = \frac{384.06}{393.6} \approx 0.98$$

Approximately 98 percent of the variation in Y is explained by variation in X.

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

TWO PROPERTIES OF r^2

1. Nonnegative quantity
2. Its limits are $0 \leq r^2 \leq 1$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

THE COEFFICIENT OF CORRELATION: R

- A measure of the degree of association between two variables

$$r = \pm \sqrt{r^2}$$


ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

PROPERTIES OF r


1. Can be positive or negative
2. Lies between the limits of -1 and 1
3. Symmetric in nature
4. Independent of the origin and scale
5. If X and Y are statistically independent, the correlation coefficient between them is zero **but zero correlation does not necessarily imply independence**
6. No meaning for describing nonlinear relations
7. Does not necessarily imply any cause and effect relationship

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

CORRELATION PATTERNS



ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)



ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

SOURCE

Gujarati, D.N. (2009) Basic Econometrics. 5th ed. Singapore, McGraw-Hill.

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)