

## Solution: Quiz 7

1. Determine every solution of the following system of linear equations by using Gaussian elimination method.

$$\begin{aligned}x - 2y &= 3z \\y + 2z - 3 &= x \\z + 3 + y &= 0\end{aligned}$$

**Solution:** By rearranging the terms in each equation, the corresponding augmented matrix of the given linear system can be written as

$$\begin{array}{l}R_1 : \\R_2 : \\R_3 : \end{array} \left[ \begin{array}{ccc|c} 1 & -2 & -3 & 0 \\ -1 & 1 & 2 & 3 \\ 0 & 1 & 1 & -3 \end{array} \right] \Leftrightarrow \begin{array}{l}x - 2y - 3z = 0 \\-x + y + 2z = 3 \\y + z = -3\end{array}$$

and we will apply the following 2 procedures in Gaussian elimination to the augmented matrix.

- I. Forward elimination to obtain the upper triangular form
- II. Back substitution

### I. Forward elimination

Step 1:  $R_1$  is the “pivot row” and  $a_{11} = 1$  is the “pivot element”.

Let  $m_{21} = \frac{a_{21}}{a_{11}} = \frac{-1}{1} = -1$ . Note that there is no need to change  $R_3$ .

$$\begin{array}{l}R_1 : \\R_2 \mapsto R_2 - m_{21}R_1 : \\R_3\end{array} \left[ \begin{array}{ccc|c} 1 & -2 & -3 & 0 \\ 0 & -1 & -1 & 3 \\ 0 & 1 & 1 & -3 \end{array} \right]$$

Step 2:  $R_2$  is the “pivot row” and  $a_{22} = -1$  is the “pivot element”.

Let  $m_{32} = \frac{a_{32}}{a_{22}} = \frac{2}{-1} = -2$ .

$$\begin{array}{l}R_1 : \\R_2 : \\R_3 \mapsto R_3 - m_{32}R_2 : \end{array} \left[ \begin{array}{ccc|c} 1 & -2 & -3 & 0 \\ 0 & -1 & -1 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right] \Leftrightarrow \begin{array}{l}x - 2y - 3z = 0 \\-y - z = 3 \\0 \cdot z = 0\end{array}$$

**II. Back substitution** Notice that any value of  $z \in \mathbb{R}$  will satisfy the last equation, where  $\mathbb{R}$  is the set of real numbers.

- Row  $R_3$ :  $z = t$ , where  $t \in \mathbb{R}$
- Row  $R_2$ :  $-y - t = 3 \Rightarrow y = -t - 3$ .
- Row  $R_1$ :  $x - 2(-t - 3) - 3t = 0 \Rightarrow x = t - 6$

That is, there are infinitely many solutions for this linear system which can be written in terms

of the set of solutions as  $\left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} t - 6 \\ -t - 3 \\ t \end{bmatrix} \mid t \in \mathbb{R} \right\}$ . ■