

Solution- Assignment 5: Part 2

Determine the partial fraction decomposition of the following rational functions.

1. $\frac{6x-1}{2x^4-x^3}$

Answer: First, we factor the denominator:

$$2x^4 - x^3 = x^3(2x - 1).$$

Since x is a repeated linear factor with the distinct linear factor $2x - 1$ in the denominator, the partial fraction decomposition is in the form

$$\frac{6x - 1}{x^3(2x - 1)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x^3} + \frac{D}{2x - 1}.$$

To find the constants A, B, C, D , consider

$$\frac{6x - 1}{x^3(2x - 1)} = \frac{Ax^2(2x - 1) + Bx(2x - 1) + C(2x - 1) + Dx^3}{x^3(2x - 1)}$$

and equate the numerators:

$$\begin{aligned} 6x - 1 &= Ax^2(2x - 1) + Bx(2x - 1) + C(2x - 1) + Dx^3 \\ &= (2A + D)x^3 + (-A + 2B)x^2 + (-B + 2C)x - C. \end{aligned}$$

First,

- set $x = 0$, we have $0 - 1 = C(0 - 1)$ which gives $C = 1$,
- set $x = 1/2$, we have $6 \cdot \frac{1}{2} - 1 = D(1/2)^3$ which gives $D = 16$.

Then we compare the coefficients of the terms with the same power of x :

$$2A + D = 0, \quad -A + 2B = 0, \quad -B + 2C = 6, \quad -C = -1.$$

Substituting $C = 1$ and $D = 16$, we have $A = -D/2 = -8$, $B = A/2 = -4$. That is, the partial fraction decomposition is

$$\frac{6x - 1}{x^3(2x - 1)} = \frac{-8}{x} + \frac{-4}{x^2} + \frac{1}{x^3} + \frac{16}{2x - 1}.$$

■

2. $\frac{x+3}{x^4+9x^2}$

Answer: First, we factor the denominator:

$$x^4 + 9x^2 = x^2(x^2 + 9).$$

Since x is a repeated linear factor with the distinct linear factor $x^2 + 9$ in the denominator, the partial fraction decomposition is in the form

$$\frac{x + 3}{x^2(x^2 + 9)} = \frac{A}{x} + \frac{B}{x^2} + \frac{Cx + D}{x^2 + 9}.$$

To find the constants A, B, C, D , consider

$$\frac{x+3}{x^2(x^2+9)} = \frac{Ax(x^2+9) + B(x^2+9) + (Cx+D)x^2}{x^2(x^2+9)}$$

and equate the numerators:

$$\begin{aligned} x+3 &= Ax(x^2+9) + B(x^2+9) + (Cx+D)x^2 \\ &= (A+C)x^3 + (B+D)x^2 + (9A)x + 9B. \end{aligned}$$

By comparing the coefficients of the terms with the same power of x :

$$A+C=0, \quad B+D=0, \quad 9A=1, \quad 9B=3.$$

That is,

$$A = \frac{1}{9}, \quad B = \frac{1}{3}, \quad C = -\frac{1}{9}, \quad D = -\frac{1}{3}.$$

Hence, the partial fraction decomposition is

$$\frac{x+3}{x^2(x^2+9)} = \frac{1}{9x} + \frac{1}{3x^2} - \frac{1}{9} \frac{x+3}{x^2+9}.$$

■

3. $\frac{x^3-2x}{x^2+3x+2}$

Answer: Since the power of the numerator is not less than the power of the denominator, we first have to divide the numerator by the denominator (e.g. long division) to obtain

$$\frac{x^3-2x}{x^2+3x+2} = x-3 + \frac{5x+6}{x^2+3x+2},$$

so that we can apply the partial fraction decomposition for $\frac{5x+6}{x^2+3x+2}$. Since $x^2+3x+2 = (x+1)(x+2)$, the decomposition is in the form

$$\begin{aligned} \frac{5x+6}{(x+1)(x+2)} &= \frac{A}{x+1} + \frac{B}{x+2} \\ \frac{5x+6}{(x+1)(x+2)} &= \frac{A(x+2) + B(x+1)}{(x+1)(x+2)} \\ 5x+6 &= A(x+2) + B(x+1) \end{aligned}$$

By setting $x = -1$, $5(-1) + 6 = A(-1 + 2)$, which gives $A = 1$ By setting $x = -2$, $5(-2) + 6 = B(-2 + 1)$, which gives $B = 4$. That is,

$$\frac{x^3-2x}{x^2+3x+2} = x-3 + \frac{1}{x+1} + \frac{4}{x+2}.$$

■

4. $\frac{x^2}{(x^2+4)}$

Answer:

Since the power of the numerator is not less than the power of the denominator, we first have to divide the numerator by the denominator (e.g. long division) to obtain

$$\frac{x^2}{(x^2+4)} = 1 - \frac{4}{x^2+4}$$

Since the divisor is an irreducible second-order polynomial and the remainder is just a constant, the remainder over the divisor, $-\frac{4}{x^2+4}$, is already in the form of partial fraction decomposition and above is the final answer. ■

5. $\frac{4x}{(x^2+1)(x^2+2x+3)}$

Answer: Since the denominator is already factored into the product of irreducible polynomials, the partial fraction decomposition is in the form

$$\frac{4x}{(x^2+1)(x^2+2x+3)} = \frac{Ax+B}{x^2+1} + \frac{Cx+D}{x^2+2x+3}$$

from which we find

$$\begin{aligned} 4x &= (Ax+B)(x^2+2x+3) + (Cx+D)(x^2+1) \\ &= (A+C)x^2 + (2A+B+D)x^2 + (3A+2B+C)x + (3B+D). \end{aligned}$$

By comparing the coefficients of the powers of x :

$$\begin{aligned} A+C &= 0 \\ 2A+B+D &= 0 \\ 3A+2B+C &= 4 \\ 3B+D &= 0 \end{aligned}$$

which give $A = 1$, $B = 1$, $C = -1$, and $D = -3$. Therefore the partial fraction decomposition is

$$\frac{4x}{(x^2+1)(x^2+2x+3)} = \frac{x+1}{x^2+1} - \frac{x+3}{x^2+2x+3}.$$

■