

**EE 325 Section 1 (Aj.Wanwiphang) Homework Assignment 1****Due date: 31 January 2020 before 11pm**

**\*\* Please submit this assignment on Moodle. For those who work on paper, please scan or submit the pictures of your work. \*\***

1. Find the answers following questions (please also show your calculation)

$$\begin{aligned} \text{a. } \sum_{i=1}^5 (a + bx_i) &= (a + bx_1) + (a + bx_2) + (a + bx_3) + (a + bx_4) + (a + bx_5) \\ &= 5a + bx_1 + bx_2 + bx_3 + bx_4 + bx_5 \\ &= 5a + b(x_1 + x_2 + x_3 + x_4 + x_5) \end{aligned}$$

$$\text{b. } \sum_{y=0}^5 f(x+y) = f(x+0) + f(x+1) + f(x+2) + f(x+3) + f(x+4) + f(x+5)$$

$$\begin{aligned} \text{c. } \sum_{i=1}^{10} i^2 &= 1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2 + 7^2 + 8^2 + 9^2 + 10^2 \\ &= 1 + 4 + 9 + 16 + 25 + 36 + 49 + 64 + 81 + 100 \\ &= 385 \end{aligned}$$

$$\begin{aligned} \text{d. } \sum_{x=1}^2 \sum_{y=2}^3 (2x+y) &= [2(1)+2] + [2(2)+3] \\ &= (2+2) + (4+3) \\ &= 4+7 = 11 \end{aligned}$$

2. Given  $X$  is discrete random variable. The probability distribution function (PDF) of this variable is shown in the table

|        |      |    |       |    |      |      |       |
|--------|------|----|-------|----|------|------|-------|
| $X$    | -2   | -1 | 0     | 1  | 2    | 3    | 4     |
| $f(x)$ | 0.5b | b  | 2.25b | 2b | 1.5b | 0.5b | 0.25b |

\*\* when b is constant number

- a. Find the value of b

$$\begin{aligned} f(x) = P(X=x) : 0.5b + b + 2.25b + 2b + 1.5b + 0.5b + 0.25b &= 1 \\ 8b &= 1 \\ b &= \frac{1}{8} = 0.125 \end{aligned}$$

- b. Find the answer for  $P(X \leq 2)$

$$\begin{aligned} P(X \leq 2) &= 1 - P(X > 2) \\ &= 1 - [P(X=3) + P(X=4)] \\ &= 1 - [0.5(0.125) + 0.25(0.125)] \end{aligned} \quad \left| \begin{aligned} &= 1 - (0.0625 + 0.03125) \\ &= 1 - 0.09375 \\ &= 0.90625 \end{aligned} \right.$$

- c. Find the answer for  $P(-2 \leq X \leq 3)$

$$\begin{aligned} P(-2 \leq X \leq 3) &= 1 - P(X=4) \\ &= 1 - 0.25(0.125) \\ &= 1 - 0.03125 = 0.96875 \end{aligned}$$


- d. Find the answer for  $P(X \geq 1)$

$$\begin{aligned} P(X \geq 1) &= P(X=1) + P(X=2) + P(X=3) + P(X=4) \\ &= 2(0.125) + 1.5(0.125) + 0.5(0.125) + 0.25(0.125) \\ &= 0.25 + 0.1875 + 0.0625 + 0.03125 \\ &= 0.53125 \end{aligned}$$

3. Given  $X$  is continuous random variable. The probability distribution function (PDF) of this variable is

$$f(x) = -\frac{1}{9}x + \frac{6}{9}, 0 \leq x \leq 3$$

- a. Plot graph for  $f(x)$



- b. Find the answer for  $P(1 \leq X \leq 3)$

$$\begin{aligned} P(1 \leq x \leq 3) &= \int_1^3 f(x) dx \\ &= \int_1^3 \left(-\frac{1}{9}x + \frac{6}{9}\right) dx \\ &= \left[-\frac{x^2}{18} + \frac{6x}{9}\right]_1^3 \\ &= \left[-\frac{3^2}{18} + \frac{6(3)}{9}\right] - \left[-\frac{1^2}{18} + \frac{6(1)}{9}\right] \\ &= \left(-\frac{9}{18} + \frac{18}{9}\right) - \left(-\frac{1}{18} + \frac{6}{9}\right) \\ &= -\frac{9}{18} + \frac{18}{9} + \frac{1}{18} - \frac{6}{9} \\ &= -\frac{8}{18} + \frac{12}{9} \\ &= -\frac{8}{18} + \frac{24}{18} \\ &= \frac{16}{18} = \frac{8}{9} \end{aligned}$$

- c. Find the answer for  $P(X \geq 2)$

$$\begin{aligned} P(x \geq 2) &= \int_2^3 f(x) dx \\ &= \int_2^3 \left(-\frac{1}{9}x + \frac{6}{9}\right) dx \\ &= \left[-\frac{x^2}{18} + \frac{6x}{9}\right]_2^3 \\ &= \left[-\frac{3^2}{18} + \frac{6(3)}{9}\right] - \left[-\frac{2^2}{18} + \frac{6(2)}{9}\right] \\ &= \left(-\frac{9}{18} + \frac{18}{9}\right) - \left(-\frac{4}{18} + \frac{12}{9}\right) \\ &= -\frac{9}{18} + \frac{18}{9} + \frac{4}{18} - \frac{12}{9} \\ &= -\frac{5}{18} + \frac{6}{9} \\ &= -\frac{5}{18} + \frac{12}{18} \\ &= \frac{7}{18} \end{aligned}$$

- d. Find the expected value of  $X$

$$\begin{aligned} E(x) &= \int_0^3 x f(x) dx \\ &= \int_0^3 x \left(-\frac{1}{9}x + \frac{6}{9}\right) dx \\ &= \int_0^3 \left(-\frac{1}{9}x^2 + \frac{6x}{9}\right) dx \\ &= \left[-\frac{1}{27}x^3 + \frac{6x^2}{18}\right]_0^3 \\ &= \left[-\frac{3^3}{27} + \frac{6(3)^2}{18}\right] - \left[-\frac{0^3}{27} + \frac{6(0)^2}{18}\right] \\ &= -1 + 3 - 0 \\ &= 2 \end{aligned}$$

4. Let random variable  $X$  be the outcome of throwing one dice and random variable  $Y$  be the outcome of tossing one coin. Coin has two sided that has valued 1 and 0.

- a. Construct the joint probability distribution function (PDF) table of  $X$  and  $Y$

| $Y \backslash X$ | 1              | 2              | 3              | 4              | 5              | 6              |
|------------------|----------------|----------------|----------------|----------------|----------------|----------------|
| 0                | $\frac{1}{12}$ | $\frac{1}{12}$ | $\frac{1}{12}$ | $\frac{1}{12}$ | $\frac{1}{12}$ | $\frac{1}{12}$ |
| 1                | $\frac{1}{12}$ | $\frac{1}{12}$ | $\frac{1}{12}$ | $\frac{1}{12}$ | $\frac{1}{12}$ | $\frac{1}{12}$ |

- b. Find the marginal probability distribution function (PDF) of  $X$

$$\begin{array}{l|l} f(X=1) = \frac{1}{6} & f(X=4) = \frac{1}{6} \\ f(X=2) = \frac{1}{6} & f(X=5) = \frac{1}{6} \\ f(X=3) = \frac{1}{6} & f(X=6) = \frac{1}{6} \end{array}$$

- c. Find the marginal probability distribution function (PDF) of  $Y$

$$\begin{array}{l} f(Y=0) = \frac{1}{2} \\ f(Y=1) = \frac{1}{2} \end{array}$$

- d. Find the conditional probability distribution function (PDF) of

$X$  given  $Y$  is equal to 1

$$\begin{aligned} f(X|Y=1) &= \frac{f(X,Y)}{f_Y(1)} \\ &= \frac{f(X,1)}{\frac{1}{2}} \end{aligned}$$

| $X$        | 1             | 2             | 3             | 4             | 5             | 6             |
|------------|---------------|---------------|---------------|---------------|---------------|---------------|
| $f(X Y=1)$ | $\frac{1}{6}$ | $\frac{1}{6}$ | $\frac{1}{6}$ | $\frac{1}{6}$ | $\frac{1}{6}$ | $\frac{1}{6}$ |

- e. Find the expected value of  $X$  given  $Y$  is equal to 1

$$\begin{aligned} E(X|Y=1) &= \frac{\sum x_i P(X=x_i|Y=1)}{P(Y=1)} = \frac{[(1 \cdot \frac{1}{12}) + (2 \cdot \frac{1}{12}) + (3 \cdot \frac{1}{12}) + (4 \cdot \frac{1}{12}) + (5 \cdot \frac{1}{12}) + (6 \cdot \frac{1}{12})]}{\frac{1}{2}} \\ &= \frac{\sum x_i P(X=x_i, Y=1)}{P(Y=1)} = \frac{7}{2} = 3.5 \end{aligned}$$

- f. Find the variance of  $X$  given  $Y$  is equal to 1

$$\begin{aligned} \text{Var}(X|Y=1) &= \sum (x - E(X|Y=1))^2 \cdot P(X|Y=1) \\ &= \left\{ \left[ \left(1 - \frac{7}{2}\right)^2 \cdot \frac{1}{6} \right] + \left[ \left(2 - \frac{7}{2}\right)^2 \cdot \frac{1}{6} \right] + \left[ \left(3 - \frac{7}{2}\right)^2 \cdot \frac{1}{6} \right] + \left[ \left(4 - \frac{7}{2}\right)^2 \cdot \frac{1}{6} \right] + \left[ \left(5 - \frac{7}{2}\right)^2 \cdot \frac{1}{6} \right] + \left[ \left(6 - \frac{7}{2}\right)^2 \cdot \frac{1}{6} \right] \right\} \\ &= \left[ \left( \frac{25}{4} \cdot \frac{1}{6} \right) + \left( \frac{9}{4} \cdot \frac{1}{6} \right) + \left( \frac{1}{4} \cdot \frac{1}{6} \right) + \left( \frac{1}{4} \cdot \frac{1}{6} \right) + \left( \frac{9}{4} \cdot \frac{1}{6} \right) + \left( \frac{25}{4} \cdot \frac{1}{6} \right) \right] \\ &= \frac{25}{24} + \frac{9}{24} + \frac{1}{24} + \frac{1}{24} + \frac{9}{24} + \frac{25}{24} \\ &= \frac{70}{24} = 2.917 \end{aligned}$$

5. If  $X_1, X_2, X_3$  is a random sample from a population with mean  $\mu$  and variance  $\sigma^2$ .  $X_1, X_2, X_3$  are not independent

$$\text{Cov}(X_1, X_2) = \text{Cov}(X_1, X_3) = \text{Cov}(X_2, X_3) = \frac{1}{4}\sigma^2$$

$\bar{X}$  is estimator used to estimate mean value.  $\bar{X} = \frac{1}{3}(X_1 + X_2 + X_3)$

Find  $E(\bar{X})$  and  $\text{var}(\bar{X})$

|                                                                                                                                                                                                                                                                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $  \begin{aligned}  E(\bar{X}) &= E\left(\frac{1}{N} \sum_{i=1}^3 X_i\right) \\  &= \frac{1}{N} E(X_1 + X_2 + X_3) \\  &= \frac{1}{N} (E(X_1) + E(X_2) + E(X_3)) \\  &= \frac{1}{3} (\mu_x + \mu_x + \mu_x) \\  &= \frac{1}{3} (3\mu_x) \\  &= \mu_x  \end{aligned}  $ | $  \begin{aligned}  \text{var}(\bar{X}) &= \text{var}\left(\frac{1}{N} \sum_{i=1}^3 X_i\right) \\  &= \frac{1}{N^2} \text{var}(X_1 + X_2 + X_3) \\  &= \frac{1}{N^2} [\text{var}(X_1) + \text{var}(X_2) + \text{var}(X_3) + \\  &\quad 2\text{cov}(X_1, X_2) + 2\text{cov}(X_1, X_3) + 2\text{cov}(X_2, X_3)] \\  &= \frac{1}{3^2} \left( \frac{1}{4} \sigma_x^2 + \frac{1}{4} \sigma_x^2 + \frac{1}{4} \sigma_x^2 + \frac{1}{2} \sigma_x^2 + \frac{1}{2} \sigma_x^2 + \frac{1}{2} \sigma_x^2 \right) \\  &= \frac{1}{9} \left( \frac{9}{4} \sigma_x^2 \right) \\  &= \frac{1}{4} \sigma_x^2 \\  &= 0.25 \sigma_x^2  \end{aligned}  $ |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

6. Given  $X_1, X_2, X_3, X_4$  are independent identically distributed random variables from population with mean  $\mu$  and variance  $\sigma^2$ .  $\bar{X}$  is estimator used to estimate mean value.  $\bar{X} = \frac{1}{4}(X_1 + X_2 + X_3 + X_4)$

- a. Find  $E(\bar{X})$  and  $\text{var}(\bar{X})$  in term of  $\mu$  and  $\sigma$

|                                                                                                                                                                                                                                                      |                                                                            |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|
| $  \begin{aligned}  E(\bar{X}) &= E\left(\frac{1}{N} \sum_{i=1}^4 X_i\right) \\  &= \frac{1}{N} E(X_1 + X_2 + X_3 + X_4) \\  &= \frac{1}{N} (E(X_1) + E(X_2) + E(X_3) + E(X_4)) \\  &= \frac{1}{4} (\mu_x + \mu_x + \mu_x + \mu_x)  \end{aligned}  $ | $  \begin{aligned}  &= \frac{1}{4} (4\mu_x) \\  &= \mu_x  \end{aligned}  $ |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|

$$\begin{aligned}
\text{var}(\bar{x}) &= \text{var}\left(\frac{1}{N} \sum_{i=1}^4 x_i\right) \\
&= \frac{1}{N^2} \text{var}(x_1 + x_2 + x_3 + x_4) \\
&= \frac{1}{4^2} (\text{var}(x_1) + \text{var}(x_2) + \text{var}(x_3) + \text{var}(x_4)) \\
&= \frac{1}{16} (6_x^2 + 6_x^2 + 6_x^2 + 6_x^2) \\
&= \frac{1}{16} (4 \cdot 6_x^2) \\
&= \frac{1}{4} 6_x^2
\end{aligned}$$

- b. Given  $\tilde{X} = \frac{1}{8}X_1 + \frac{1}{4}X_2 + \frac{1}{8}X_3 + \frac{1}{2}X_4$  is another estimator of  $\mu$ . Show that  $\tilde{X}$  is an unbiased estimator of  $\mu$

$$\begin{aligned}
\tilde{X} &= \frac{1}{8}x_1 + \frac{1}{4}x_2 + \frac{1}{8}x_3 + \frac{1}{2}x_4 \\
&= \frac{1}{4} (0.5x_1 + x_2 + 0.5x_3 + 2x_4) \\
E(\tilde{x}) &= E\left(\frac{1}{N} \sum_{i=1}^4 x_i\right) \\
&= \frac{1}{4} E(0.5x_1 + x_2 + 0.5x_3 + 2x_4) \\
&= \frac{1}{4} [E(0.5x_1) + E(x_2) + E(0.5x_3) + E(2x_4)] \\
&= \frac{1}{4} [0.5E(x_1) + E(x_2) + 0.5E(x_3) + 2E(x_4)] \\
&= \frac{1}{4} (0.5\mu_x + \mu_x + 0.5\mu_x + 2\mu_x) \\
&= \frac{1}{4} (4\mu_x) \\
&= \mu_x
\end{aligned}$$

So,  $\tilde{x}$  is an unbiased estimator of  $\mu$

- c. Between  $\bar{X}$  and  $\tilde{X}$ , which one is the better estimator for  $\mu$ ? Why?

Between  $\bar{x}$  and  $\tilde{x}$ ,  $\bar{x}$  is the better estimator for  $\mu$  than  $\tilde{x}$  because  $\bar{x}$  has smaller value of variance than  $\tilde{x}$  which is  $\bar{x} = 0.25 \cdot 6_x^2 < \tilde{x} = 0.34 \cdot 6_x^2$