

TWO-VARIABLE REGRESSION: INTERVAL ESTIMATION AND HYPOTHESIS TESTING

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

The theory of estimation consists of

- **Point estimation**

A single value used to approximate a population parameter

- **Interval estimation**

a range of values estimating the parameter (may or may not contain the value of the parameter being estimated)

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

Confidence level of an interval estimate of a parameter:

the probability that the interval estimate will contain the parameter, assuming that a large number of samples are selected and that the estimation process on the same parameter is repeated

E.g. 90% Confidence level
95% Confidence level
99% Confidence level

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

INTERVAL ESTIMATION

- We want to find how “close”, say, $\hat{\beta}_2$ is to β_2
- We try to find out δ and α such that the probability that the **random interval** contains the true β_2 is $(\hat{\beta}_2 - \delta, \hat{\beta}_2 + \delta)$
- Confidence interval is $1 - \alpha$

$$\Pr(\hat{\beta}_2 - \delta \leq \beta_2 \leq \hat{\beta}_2 + \delta) = 1 - \alpha$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

INTERVAL ESTIMATION

- $1 - \alpha$ is known as confidence coefficient
- α is known as the level of significance

E.g. 95% confidence level
 $\alpha = 1 - 0.95 = 0.05$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

α is level of significance

- The probability of committing a type I error
- A type I error consists in rejecting a true hypothesis

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\alpha = 0.05$$

$$t_{\alpha/2, df=10-2} =$$

$$t_{\alpha, df=10-2} =$$

$$t_{\alpha/2, df=200-2} =$$

$$t_{\alpha, df=200-2} =$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\alpha = 0.01$$

$$t_{\alpha/2, df=10-2} =$$

$$t_{\alpha, df=10-2} =$$

$$t_{\alpha/2, df=200-2} =$$

$$t_{\alpha, df=200-2} =$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

CONFIDENCE INTERVALS FOR REGRESSION COEFFICIENTS β_1 AND β_2

- With the normality assumption for u_i
- The OLS estimators $\hat{\beta}_1$ and $\hat{\beta}_2$ are themselves normally distributed with means and variances
- Variance (σ^2) is known

$$\begin{aligned} Z &= \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)} \\ &= \frac{(\hat{\beta}_2 - \beta_2) \sqrt{\sum x_i^2}}{\sigma} \end{aligned}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

CONFIDENCE INTERVALS FOR REGRESSION COEFFICIENTS β_1 AND β_2

- σ^2 is rarely known, in practice it is determined by the unbiased estimator $\hat{\sigma}^2$

$$\begin{aligned} t &= \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)} \\ &= \frac{(\hat{\beta}_2 - \beta_2) \sqrt{\sum x_i^2}}{\hat{\sigma}} \end{aligned}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

CONFIDENCE INTERVALS FOR REGRESSION COEFFICIENTS β_1 AND β_2

$$\Pr(-t_{\alpha/2} \leq t \leq t_{\alpha/2}) = 1 - \alpha$$

$$\Pr[-t_{\alpha/2} \leq \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)} \leq t_{\alpha/2}] = 1 - \alpha$$

$$\Pr[\hat{\beta}_2 - t_{\alpha/2} se(\hat{\beta}_2) \leq \beta_2 \leq \hat{\beta}_2 + t_{\alpha/2} se(\hat{\beta}_2)] = 1 - \alpha$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

CONFIDENCE INTERVALS FOR REGRESSION COEFFICIENTS β_2 AND

100(1- α)% confidence interval for β_2 :

$$\hat{\beta}_2 \pm t_{\alpha/2} se(\hat{\beta}_2)$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

TABLE 2.6
Mean Hourly Wage
by Education

Source: Arthur S.
Goldberger, *Introductory
Econometrics*, Harvard
University Press, Cambridge,
Mass., 1998, Table 1.1, p. 5
(adapted).

Years of Schooling	Mean Wage, \$	Number of People
6	4.4567	3
7	5.7700	5
8	5.9787	15
9	7.3317	12
10	7.3182	17
11	6.5844	27
12	7.8182	218
13	7.8351	37
14	11.0223	56
15	10.6738	13
16	10.8361	70
17	13.6150	24
18	13.5310	31
Total		528

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

CONFIDENCE INTERVALS FOR REGRESSION COEFFICIENTS β_1 AND β_2

Example: Mean hourly wages (Y) on education (X)

$$\hat{\beta}_2 = 0.7240$$

$$se(\hat{\beta}_2) = 0.0700$$

$$n = 13, df = n - 2 = 11, \alpha = 0.05$$

$$1 - \alpha = 0.95$$

$$\text{critical } t_{\alpha/2} = 2.201$$

$$\Pr[\hat{\beta}_2 - t_{\alpha/2} se(\hat{\beta}_2) \leq \beta_2 \leq \hat{\beta}_2 + t_{\alpha/2} se(\hat{\beta}_2)] = 1 - \alpha$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

CONFIDENCE INTERVALS FOR REGRESSION COEFFICIENTS β_1 AND β_2

$$0.5700 \leq \beta_2 \leq 0.8780$$

or

$$0.7240 \pm 2.201(0.0700)$$

Given the confidence coefficient of 95 percent, the true parameter is between 0.5700 and 0.8780.

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

CONFIDENCE INTERVALS FOR REGRESSION COEFFICIENTS β_1 AND β_2

The **correct interpretation** of this confidence interval is:

Given the confidence coefficient of 95 percent, in 95 out of 100 cases intervals will contain the true β_2

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

CONFIDENCE INTERVALS FOR REGRESSION COEFFICIENTS β_1 AND β_2

The **wrong interpretation** of this confidence interval is:

The probability is 95 percent that the specific interval contains the true

Since the interval is now fixed and no longer random; therefore β_2 either lies in it or it does not β_2

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

CLASS EXERCISE

Example: Mean hourly wages (Y) on education (X)

$$\hat{\beta}_2 = 0.7240$$

$$se(\hat{\beta}_2) = 0.0700$$

$$n = 13, df = n - 2 = 11, \alpha = 0.01$$

$$1 - \alpha = 0.99$$

$$\text{critical } t_{\alpha/2} =$$

$$\Pr[\hat{\beta}_2 - t_{\alpha/2} se(\hat{\beta}_2) \leq \beta_2 \leq \hat{\beta}_2 + t_{\alpha/2} se(\hat{\beta}_2)] = 1 - \alpha$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)


CONFIDENCE INTERVALS FOR σ^2

Under the normality assumption, the variable follows the χ^2 distribution with $n-2$ df

$$\chi^2 = (n-2) \frac{\hat{\sigma}^2}{\sigma^2}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

CONFIDENCE INTERVALS FOR σ^2



ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\alpha = 0.05$$

$$\chi_{1-\alpha/2, df=10-2} =$$

$$\chi_{\alpha/2, df=10-2} =$$

$$\chi_{1-\alpha/2, df=200-2} =$$

$$\chi_{\alpha/2, df=200-2} =$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\alpha = 0.01$$

$$\chi_{1-\alpha/2, df=10-2} =$$

$$\chi_{\alpha/2, df=10-2} =$$

$$\chi_{1-\alpha/2, df=200-2} =$$

$$\chi_{\alpha/2, df=200-2} =$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

CONFIDENCE INTERVALS FOR σ^2

$$\Pr(\chi_{1-\alpha/2}^2 \leq \chi^2 \leq \chi_{\alpha/2}^2) = 1 - \alpha$$

$$\Pr\left[(n-2) \frac{\hat{\sigma}^2}{\chi_{\alpha/2}^2} \leq \sigma^2 \leq (n-2) \frac{\hat{\sigma}^2}{\chi_{1-\alpha/2}^2}\right] = 1 - \alpha$$

Which gives the $100(1-\alpha)\%$ confidence interval for σ^2

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

CONFIDENCE INTERVALS FOR σ^2

Example: Wage-education $\hat{\sigma}^2 = 0.8936$

$$\alpha = 0.05, df = n - 2 = 11$$

$$\chi_{0.025}^2 = 21.9200, \chi_{0.975}^2 = 3.8157$$

$$0.4484 \leq \sigma^2 \leq 2.5760$$

Given the confidence coefficient of 95 percent, the true parameter is between 0.4484 and 2.5760

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

CONFIDENCE INTERVALS FOR σ^2

The interpretation of this interval is:

If we established 95 percent confidence limits on σ^2

And if we maintain a priori that these limits will

Include the true σ^2 , we will be right in the long run 95 percent of the time

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

CLASS EXERCISE

Example: Wage-

education $\hat{\sigma}^2 = 0.8936$

$$\alpha = 0.01, df = n - 2 = 11$$

$$\chi^2_{0.025} = \underline{\hspace{2cm}}, \chi^2_{0.975} = \underline{\hspace{2cm}}$$

$$\underline{\hspace{2cm}} \leq \sigma^2 \leq \underline{\hspace{2cm}}$$

Given the confidence coefficient of 99 percent, the true parameter is between and

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

HYPOTHESIS TESTING

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

HYPOTHESIS TESTING

In statistics, a **hypothesis** is a claim or statement about a property of a population.

A **hypothesis test** (or **test of significance**) is a standard procedure for testing a claim about a property of a population.

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

HYPOTHESIS TESTING

1. Define population in study
2. State the hypothesis to be investigated
3. Give the desired significance level
4. Select a sample from population
5. Collect the data
6. Perform the calculations
7. Reach a conclusion

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

HYPOTHESIS TESTING

- **Null Hypothesis (H_0):**

A Statistical hypothesis stating there is no difference between a parameter and a specific value

- We test the null hypothesis directly
- Either reject H_0 or fail to reject H_0

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

HYPOTHESIS TESTING

- **Alternative hypothesis:** (H_1):
Stating the existence of a difference between a parameter and a specific value.
- The symbolic form of the alternative hypothesis must use one of these symbols: $\neq$, $<$, $>$.

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

TABLE 5.1
The t Test of
Significance: Decision
Rules

Type of Hypothesis	H_0 : The Null Hypothesis	H_1 : The Alternative Hypothesis	Decision Rule: Reject H_0 If
Two-tail	$\beta_2 = \beta_2^0$	$\beta_2 \neq \beta_2^0$	$ t > t_{\alpha/2, df}$
Right-tail	$\beta_2 \leq \beta_2^0$	$\beta_2 > \beta_2^0$	$t > t_{\alpha, df}$
Left-tail	$\beta_2 \geq \beta_2^0$	$\beta_2 < \beta_2^0$	$t < -t_{\alpha, df}$

Notes: β_2^0 is the hypothesized numerical value of β_2 .
 $|t|$ means the absolute value of t .
 t_{α} or $t_{\alpha/2}$ means the critical t value at the α or $\alpha/2$ level of significance.
 df : degrees of freedom, $(n - 2)$ for the two-variable model, $(n - 3)$ for the three-variable model, and so on.
The same procedure holds to test hypotheses about β_1 .

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

HYPOTHESIS TESTING: THE CONFIDENCE-INTERVAL APPROACH

Two-Sided or Two-Tail Test

$$H_0: \beta_2 = 0.5$$

$$H_1: \beta_2 \neq 0.5$$

Therefore, if $\hat{\beta}_2$ under H_0 falls within the $100(1-\alpha)\%$ Confidence interval, we do not reject the null hypothesis; if it lies outside the interval, we may reject it

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\Pr[-t_{\alpha/2} \leq \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)} \leq t_{\alpha/2}] = 1 - \alpha$$

$-t_{\alpha/2}$ and $t_{\alpha/2}$ are the value of t (the critical t value)

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

THE TEST-OF-SIGNIFICANCE OF REGRESSION COEFFICIENTS: THE T TEST

$$\Pr[\hat{\beta}_2 - t_{\alpha/2} se(\hat{\beta}_2) \leq \beta_2 \leq \hat{\beta}_2 + t_{\alpha/2} se(\hat{\beta}_2)] = 1 - \alpha$$

Region of acceptance


The $100(1-\alpha)\%$ confidence interval

Region of rejection

The regions outside the confidence interval

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

HYPOTHESIS TESTING: THE CONFIDENCE-INTERVAL APPROACH



ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

HYPOTHESIS TESTING: THE CONFIDENCE-INTERVAL APPROACH

- When we reject the null hypothesis, we say that our finding is **statistically significant**
- When we do not reject the null hypothesis, we say that our finding is **not statistically significant**

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

In the **confidence-interval procedure** we try to establish a range or an interval that has a certain probability of including the true but unknown

, whereas in the test-of-significance approach we hypothesize some value for β_2 and try to see whether the computed $\hat{\beta}_2$ lies within reasonable (confidence) limits around the hypothesized value

$$\hat{\beta}_2$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

THE TEST-OF-SIGNIFICANCE OF REGRESSION COEFFICIENTS: THE T TEST

Under the normality assumption the variable

$$t = \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)} = \frac{(\hat{\beta}_2 - \beta_2)\sqrt{\sum x_i^2}}{\hat{\sigma}} = \frac{(\hat{\beta}_2 - \beta_2)\sqrt{\sum (X_i - \bar{X})^2}}{\hat{\sigma}}$$

follows the t distribution with n-2 df

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

THE TEST-OF-SIGNIFICANCE OF REGRESSION COEFFICIENTS: THE T TEST

Example: Wage-education

$$\hat{\beta}_2 = 0.7240$$

$$se(\hat{\beta}_2) = 0.0700$$

$$df = 11, \alpha = 0.05, t_{\alpha/2} = 2.201$$

$$H_0 : \beta_2 = 0.5$$


$$H_1 : \beta_2 \neq 0.5$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

THE TEST-OF-SIGNIFICANCE OF REGRESSION COEFFICIENTS: THE T TEST

$$\Pr[-t_{\alpha/2} \leq \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)} \leq t_{\alpha/2}] = 1 - \alpha$$


$$t = \frac{0.7240 - 0.5}{0.0700} = 3.2$$



Reject H_0

THE TEST-OF-SIGNIFICANCE OF REGRESSION COEFFICIENTS: THE T TEST

$$t = \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)} = \frac{(\hat{\beta}_2 - \beta_2)\sqrt{\sum x_i^2}}{\hat{\sigma}} = \frac{(\hat{\beta}_2 - \beta_2)\sqrt{\sum (X_i - \bar{X})^2}}{\hat{\sigma}}$$



H_0
Reject

Tangtipongkul)

THE TEST-OF-SIGNIFICANCE OF REGRESSION COEFFICIENTS: THE T TEST

Significance tests

A statistic is said to be statistically significant if the value of the test statistic **lies in the critical region**



The null hypothesis is rejected

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

ONE TAILED TEST

Y Sales	X Price
49	1
45	2
44	3
39	4
38	5
37	6
34	7
33	8
30	9
29	10

$$\hat{Y}_i = 49.667 - 2.1576X_i$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

RIGHT TAIL TEST

$$\hat{\beta}_2 = -2.1576$$

$$se(\hat{\beta}_2) = 0.1204$$

$$df = 10 - 2 = 8, \alpha = 0.05, t_{0.05} = 1.860$$

$$H_0 : \beta_2 \leq 0$$

$$H_1 : \beta_2 > 0$$

$$t = \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)} = \frac{-2.1576 - 0}{0.1204} = -17.92$$

Not reject H_0 . There is not enough evidence to say that

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

LEFT TAIL TEST

$$\hat{\beta}_2 = -2.1576$$

$$se(\hat{\beta}_2) = 0.1204$$

$$df = 10 - 2 = 8, \alpha = 0.05, t_{0.05} = -1.860$$

$$H_0 : \beta_2 \geq 0$$

$$H_1 : \beta_2 < 0$$

$$t = \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)} = \frac{-2.1576 - 0}{0.1204} = -17.92$$

Reject H_0 . There is enough evidence to say that

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

DECISION RULE:

- Right tailed test
If $t > \text{critical } (t_\alpha)$ reject H_0
- Left tailed test
If $t < -\text{critical } (-t_\alpha)$ reject H_0

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

EXTRA CREDIT

- The Confidence interval approach

$$\Pr[\hat{\beta}_2 - t_{\alpha/2} se(\hat{\beta}_2) \leq \beta_2 \leq \hat{\beta}_2 + t_{\alpha/2} se(\hat{\beta}_2)] = 1 - \alpha$$

- The Test of significance approach

$$t = \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)} = \frac{(\hat{\beta}_2 - \beta_2) \sqrt{\sum x_i^2}}{\hat{\sigma}} = \frac{(\hat{\beta}_2 - \beta_2) \sqrt{\sum (X_i - \bar{X})^2}}{\hat{\sigma}}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

HYPOTHESIS TESTING

Example: Wage-education

$$\hat{\beta}_2 = 0.7240$$

$$se(\hat{\beta}_2) = 0.0700$$

$$df = 11, \alpha = 0.01, t_{\alpha/2} =$$

Step 1

$$H_0 : \beta_2 = 0$$

$$H_1 : \beta_2 \neq 0$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

THE CONFIDENCE INTERVAL APPROACH

$$\Pr[\hat{\beta}_2 - t_{\alpha/2} se(\hat{\beta}_2) \leq \beta_2 \leq \hat{\beta}_2 + t_{\alpha/2} se(\hat{\beta}_2)] = 1 - \alpha$$

$$\Pr[0.7240 - 3.106(0.0700) \leq \beta_2 \leq 0.7240 + 3.106(0.0700)] = 1 - 0.01$$

$$\Pr[0.5066 \leq \beta_2 \leq 0.9414] = 0.99$$

The confidence interval doesn't contain zero.

Reject H_0

There is enough evidence to say that $\beta_2 \neq 0$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

THE TEST OF SIGNIFICANCE APPROACH

$$t = \frac{\hat{\beta}_2 - \beta_2}{se(\hat{\beta}_2)}$$

$$= \frac{0.7240 - 0}{0.0700}$$

$$= 10.3429$$

Critical t value = ± 3.106

$t > \text{Critical } t \text{ value}$

Reject H_0

There is enough evidence to say that $\beta_2 \neq 0$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

THE TEST-OF-SIGNIFICANCE OF σ^2 : THE χ^2 TEST

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

THE TEST-OF-SIGNIFICANCE OF σ^2 : THE χ^2 TEST

$$\chi^2 = (n-2) \frac{\hat{\sigma}^2}{\sigma^2}$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

TABLE 5.2
A Summary of the
 χ^2 Test

H_0 : The Null Hypothesis	H_1 : The Alternative Hypothesis	Critical Region: Reject H_0 if
$\sigma^2 = \sigma_0^2$	$\sigma^2 > \sigma_0^2$	$\frac{df(\hat{\sigma}^2)}{\sigma_0^2} > \chi_{\alpha, df}^2$
$\sigma^2 = \sigma_0^2$	$\sigma^2 < \sigma_0^2$	$\frac{df(\hat{\sigma}^2)}{\sigma_0^2} < \chi_{(1-\alpha), df}^2$
$\sigma^2 = \sigma_0^2$	$\sigma^2 \neq \sigma_0^2$	$\frac{df(\hat{\sigma}^2)}{\sigma_0^2} > \chi_{\alpha/2, df}^2$ or $< \chi_{(1-\alpha/2), df}^2$

Note: σ_0^2 is the value of σ^2 under the null hypothesis. The first subscript on χ^2 in the last column is the level of significance, and the second subscript is the degrees of freedom. These are critical chi-square values. Note that df is $(n-2)$ for the two-variable regression model, $(n-3)$ for the three-variable regression model, and so on.

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

THE TEST-OF-SIGNIFICANCE OF σ^2 : THE χ^2 TEST

example

$$\hat{\sigma}^2 = 0.8937, df = 11$$

$$H_0: \sigma^2 = 0.6$$

$$H_1: \sigma^2 \neq 0.6$$

$$\chi^2 = (n-2) \frac{\hat{\sigma}^2}{\sigma^2} = 16.3845$$

$\alpha = 0.05$, the critical χ^2 values are $\chi^2_{1-\alpha/2} = 3.81575$ and $\chi^2_{\alpha/2} = 21.9200$


We do not reject the null hypothesis.

There is not enough evidence to say that σ^2 is different from 0.6.

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

ANALYSIS OF VARIANCE

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)



ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$TSS = ESS + RSS$$

$$\sum y_i^2 = \sum (Y_i - \bar{Y})^2 \quad (\text{Total Sum of Squares, TSS})$$

$$\sum \hat{y}_i^2 = \sum (\hat{Y}_i - \bar{Y})^2 = \hat{\beta}_2^2 \sum x_i^2$$

(Explained Sum of Squares, ESS)

$$\sum \hat{u}_i^2$$

(Residual Sum of Squares, RSS)

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

ANALYSIS OF VARIANCE

$$\sum y_i^2 = \sum \hat{y}_i^2 + \sum \hat{u}_i^2 = \hat{\beta}_2^2 \sum x_i^2 + \sum \hat{u}_i^2$$

$$TSS = ESS + RSS$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

ANALYSIS OF VARIANCE

TABLE 5.3
ANOVA Table for the
Two-Variable
Regression Model

Source of Variation	SS*	df	MSS†
Due to regression (ESS)	$\sum \hat{y}_i^2 = \hat{\beta}_2^2 \sum x_i^2$	1	$\hat{\beta}_2^2 \sum x_i^2$
Due to residuals (RSS)	$\sum \hat{u}_i^2$	$n - 2$	$\frac{\sum \hat{u}_i^2}{n - 2} = \hat{\sigma}^2$
TSS	$\sum y_i^2$	$n - 1$	

*SS means sum of squares.

†Mean sum of squares, which is obtained by dividing SS by their df.

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

ANALYSIS OF VARIANCE

$$F = \frac{MSS \text{ of ESS}}{MSS \text{ of RSS}}$$

$$= \frac{\hat{\beta}_2^2 \sum x_i^2}{\sum \hat{u}_i^2 / (n-2)}$$

$$= \frac{\hat{\beta}_2^2 \sum x_i^2}{\hat{\sigma}^2}$$

F distribution with 1 df in the numerator and
(n-2) df in the denominator

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

MSE (Mean of Explained Sum of Square)

MSR (Mean of Residual Sum of Square)

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

$$\alpha = 0.05$$

$$F_{\alpha,1,5} =$$

$$F_{\alpha,1,8} =$$

$$F_{\alpha,2,5} =$$

$$F_{\alpha,3,5} =$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

• If $F^* > \text{Critical F value } (F_{\alpha;1,n-2})$ Reject H_0

• If $F^* < \text{Critical F value } (F_{\alpha;1,n-2})$ Not reject

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

$$\alpha = 0.01$$

$$F_{\alpha,1,5} =$$

$$F_{\alpha,1,8} =$$

$$F_{\alpha,2,5} =$$

$$F_{\alpha,3,5} =$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

Example: Wage-Education

TABLE 2.6
Mean Hourly Wage
by Education

Source: Arthur S.
Goldberger, *Introductory
Econometrics*, Harvard
University Press, Cambridge,
Mass., 1998, Table 1.1, p. 5
(adapted).

Years of Schooling	Mean Wage, \$	Number of People
6	4.4567	3
7	5.7700	5
8	5.9787	15
9	7.3317	12
10	7.3182	17
11	6.5844	27
12	7.8182	218
13	7.8351	37
14	11.0223	56
15	10.6738	13
16	10.8361	70
17	13.6150	24
18	13.5310	31
		Total 528

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

F-Test

$$H_0 : \beta_2 = 0$$

$$H_1 : \beta_2 \neq 0$$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

TABLE 5.4

ANOVA Table for the
Wages-Education
Example

Source of Variation	SS	df	MSS	
Due to regression (ESS)	95.4255	1	95.4255	$F = \frac{95.4255}{0.8811}$
Due to residuals (RSS)	9.6928	11	0.8811	$= 108.3026$
TSS	105.1183	12		

If $F^* > \text{Critical F value } (F_{0.05;1,11})$ Reject H_0

If $F^* < \text{Critical F value } (F_{0.05;1,11})$ Not reject H_0

$(F_{0.05;1,11}) = 4.84$

Reject

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

APPLICATION OF REGRESSION ANALYSIS: THE PROBLEM OF PREDICTION

• Mean prediction

- Prediction of the conditional mean value of Y corresponding to a chosen X, say X_0 , that is the point on the population regression line itself

• Individual prediction

- Prediction of an individual Y value corresponding to X_0

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

TABLE 2.6

Mean Hourly Wage
by Education

Source: Arthur S.
Goldberger, *Introductory
Econometrics*, Harvard
University Press, Cambridge,
Mass., 1998, Table 1.1, p. 5
(adapted).

Years of Schooling	Mean Wage, \$	Number of People
6	4.4567	3
7	5.7700	5
8	5.9787	15
9	7.3317	12
10	7.3182	17
11	6.5844	27
12	7.8182	218
13	7.8351	37
14	11.0223	56
15	10.6738	13
16	10.8361	70
17	13.6150	24
18	13.5310	31
		Total 528

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

MEAN PREDICTION

$$\hat{Y}_i = -0.0144 + 0.7240 X_i$$

Assume that $X_0 = 20$

$$E(Y | X_0 = 20) = ?$$

$$\begin{aligned}\hat{Y}_0 &= -0.0144 + 0.7240 X_0 \\ &= -0.0144 + 0.7240(20) \\ &= 14.4656\end{aligned}$$

where $\hat{Y}_0$ = estimator of $E(Y | X_0)$

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

MEAN PREDICTION

$\hat{Y}_0$ is normally distributed with mean $(\beta_1 + \beta_2 X_0)$ and

$$\text{var}(\hat{Y}_0) = \sigma^2 \left[\frac{1}{n} + \frac{(X_0 - \bar{X})^2}{\sum x_i^2} \right]$$

By replace the unknown σ^2 by its unbiased estimators $\hat{\sigma}^2$

$$t = \frac{\hat{Y}_0 - (\beta_1 + \beta_2 X_0)}{\text{se}(\hat{Y}_0)}$$

follows the t distribution with n-2 df

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)

MEAN PREDICTION

The t distribution can therefore be used to derive confidence intervals for the true $E(Y_0 | X_0)$

$$\Pr[\hat{\beta}_1 + \hat{\beta}_2 X_0 - t_{\alpha/2} se(\hat{Y}_0) \leq \beta_1 + \beta_2 X_0 \leq \hat{\beta}_1 + \hat{\beta}_2 X_0 + t_{\alpha/2} se(\hat{Y}_0)] = 1 - \alpha$$

where $se(\hat{Y}_0)$ is obtained from

$$\text{var}(\hat{Y}_0) = \sigma^2 \left[\frac{1}{n} + \frac{(X_0 - \bar{X})^2}{\sum X_i^2} \right]$$

$$\text{var}(\hat{Y}_0) = 0.8936 \left[\frac{1}{13} + \frac{(20 - 12)^2}{182} \right] = 0.3826$$

$$se(\hat{Y}_0) = 0.6185$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

MEAN PREDICTION


Therefore, the 95 percent confidence interval for true

$E(Y | X_0) = \beta_1 + \beta_2 X_0$ is given by

$$14.4656 - 2.201(0.6185) \leq E(Y_0 | X = 20) \leq 14.4656 + 2.20(0.6185)$$

$$13.1043 \leq E(Y_0 | X = 20) \leq 15.8260$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)



ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

INDIVIDUAL PREDICTION

$$t = \frac{Y_0 - \hat{Y}_0}{se(Y_0 - \hat{Y}_0)}$$

$$\text{var}(Y_0 - \hat{Y}_0) = E[Y_0 - \hat{Y}_0]^2 = \sigma^2 \left[1 + \frac{1}{n} + \frac{(X_0 - \bar{X})^2}{\sum X_i^2} \right]$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

$$\hat{Y}_0 \pm t_{\alpha/2} \hat{\sigma}\{Y_0 - \hat{Y}_0\}$$

$$\hat{\sigma}\{Y_0 - \hat{Y}_0\} = \hat{\sigma} \sqrt{1 + \frac{1}{n} + \frac{(X_0 - \bar{X})^2}{\sum (X_i - \bar{X})^2}}$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

Therefore, the t distribution can be used to draw inferences about the true Y_0 . The point prediction of Y_0 is 14.4656, the same as that of Y_0 , and its variance is 1.2357

$$(12.0190 \leq Y_0 | X_0 = 20 \leq 16.9122)$$

ee325 2/2012 (Ajarn Kaewkwan Tangtipongkul)

SOURCE

Gujarati, D.N. (2009) Basic Econometrics. 5th ed. Singapore, McGraw-Hill. (G)

ee325 2/2012 (Ajarn Kaewkwan
Tangtipongkul)